

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 April 2001 (26.04.2001)

PCT

(10) International Publication Number
WO 01/29011 A3

(51) International Patent Classification: C07D 243/12,
A61K 31/55, A61P 43/00, C07D 409/04, 405/04, 417/04,
401/04, 403/10, 401/14, 417/14, 413/10, 405/06, 401/06,
417/10, C07F 7/10

(21) International Application Number: PCT/EP00/09553

(22) International Filing Date:
29 September 2000 (29.09.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
99120520.4 15 October 1999 (15.10.1999) EP

(71) Applicant: F. HOFFMANN-LA ROCHE AG [CH/CH];
124 Grenzacherstrasse, CH-4070 Basel (CH).

(72) Inventors: ADAM, Geo; 8 Untere Staltenstrasse, 79650
Schopfheim (DE). ALANINE, Alexander; 5, rue d'Al-
tkirch, F-68400 Riedisheim (FR). GOETSCHL, Erwin;
23 Landhofweg, CH-4153 Reinach (CH). MUTEL, Vin-
cent; 15, place des Marechaux, F-68100 Mulhouse (FR).
WOLTERING, Thomas, Johannes; 13 Riedlistrasse,
79576 Weil am Rhein (DE).

(74) Agent: WAECHTER, Dieter; 124 Grenzacherstrasse,
CH-4070 Basle (CH).

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ,
BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE,
ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP,
KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA,
MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU,
SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ,
VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:


— with international search report

(88) Date of publication of the international search report:
8 November 2001

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

WO 01/29011 A3

(54) Title: BENZODIAZEPINE DERIVATIVES AS METABOTROPIC GLUTAMATE RECEPTOR ANTAGONISTS



(I)

(57) Abstract: The present invention relates to compounds of general formula (I) wherein X is a single bond or an ethynediyl group, wherein, in case X is a single bond, R¹ is halo- gen or phenyl which is optionally substituted with halogen, lower alkyl, halo-lower alkyl, lower alkoxy, halo-lower alkoxy, or cyano; In case X is an ethynediyl group, R¹ is phenyl, optionally substituted with halogen, lower alkyl, halo-lower alkyl, lower cycloalkyl, lower alkoxy or halo-lower alkoxy; R² is a 5 or 6 membered aryl or heteroaryl which are optionally substituted. The compounds according to the present invention can be used for treating or preventing acute and/or chronic neurological disorders such as psychosis, schizophrenia, Alzheimer's disease, cognitive disorders and memory deficits.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 00/09553

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D243/12 A61K31/55 A61P43/00 C07D409/04 C07D405/04
C07D417/04 C07D401/04 C07D403/10 C07D401/14 C07D417/14
C07D413/10 C07D405/06 C07D401/06 C07D417/10 C07F7/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P C07F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X.	M. HANUMANTHA RAO ET AL.: "Synthesis of 1H-1,5-benzodiazepin-2(3H)-ones from 5(4H)-isoxazolone, a heterocyclic bifunctional C-3 synthon" SYNTHESIS., vol. 5, - 1992 pages 446-448, XP002167739 GEORG THIEME VERLAG. STUTTGART., DE ISSN: 0039-7881 * compounds 3a-3e *	1
A	WO 97 05109 A (NOVO NORDISK A/S) 13 February 1997 (1997-02-13) claims 1,9	1,6,7
A	WO 99 26927 A (NPS PHARMACEUTICALS, INC.) 3 June 1999 (1999-06-03) page 6, line 19 - line 33; claim 1; figure 1	1,6,7

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

18 May 2001

Date of making of the international search report

01/06/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax. (+31-70) 340-3016

Authorized officer

Van Bijlen, H

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 00/09553

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9705109 A	13-02-1997	AU 6514296 A	26-02-1997
		EP 0843660 A	27-05-1998
		JP 11509847 T	31-08-1999
		US 5696148 A	09-12-1997
WO 9926927 A	03-06-1999	AU 1531799 A	15-06-1999
		CN 1285820 T	28-02-2001
		EP 1037878 A	27-09-2000

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau




(43) International Publication Date
26 April 2001 (26.04.2001)

PCT

(10) International Publication Number
WO 01/29011 A2

- (51) International Patent Classification⁷: **C07D 243/00** (74) Agent: **WAECHTER, Dieter**; 124 Grenzacherstrasse, CH-4070 Basle (CH).
- (21) International Application Number: **PCT/EP00/09553**
- (22) International Filing Date:
29 September 2000 (29.09.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
99120520.4 15 October 1999 (15.10.1999) EP
- (71) Applicant: **F. HOFFMANN-LA ROCHE AG [CH/CH]**;
124 Grenzacherstrasse, CH-4070 Basel (CH).
- (81) Designated States (*national*): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (72) Inventors: **ADAM, Geo**; 8 Untere Staltenstrasse, 79650 Schopfheim (DE). **ALANINE, Alexander**; 5, rue d'Altkirch, F-68400 Riedisheim (FR). **GOETSCHI, Erwin**; 23 Landhofweg, CH-4153 Reinach (CH). **MUTEL, Vincent**; 15, place des Marechaux, F-68100 Mulhouse (FR). **WOLTERING, Thomas, Johannes**; 13 Riedlistrasse, 79576 Weil am Rhein (DE).
- Published:**
— *Without international search report and to be republished upon receipt of that report.*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: **BENZODIAZEPINE DERIVATIVES**




(57) Abstract: The present invention relates to compounds of general formula (I) wherein X is a single bond or an ethynediyl group, wherein, in case X is a single bond, R¹ is halogen or phenyl which is optionally substituted with halogen, lower alkyl, halo-lower alkyl, lower alkoxy, halo-lower alkoxy, or cyano; In case X is an ethynediyl group, R¹ is phenyl, optionally substituted with halogen, lower alkyl, halo-lower alkyl, lower cycloalkyl, lower alkoxy or halo-lower alkoxy; R³ is a 5 or 6 membered aryl or heteroaryl which are optionally substituted. The compounds according to the present invention can be used for treating or preventing acute and/or chronic neurological disorders such as psychosis, schizophrenia, Alzheimer's disease, cognitive disorders and memory deficits.

WO 01/29011 A2

Benzodiazepine derivatives

The present invention relates to compounds of general formula I



I

wherein

- 5 X is a single bond or an ethynediyl group, wherein,

In case X is a single bond,

R¹ is halogen or phenyl which is optionally substituted with halogen, lower alkyl, halo-lower alkyl, lower alkoxy, halo-lower alkoxy, or cyano;

In case X is an ethynediyl group,

- 10 R¹ is phenyl, optionally substituted with halogen, lower alkyl, halo-lower alkyl, lower cycloalkyl, lower alkoxy or halo-lower alkoxy;

R² is halogen; hydroxy; lower alkyl; lower halo-alkyl; lower alkoxy; hydroxymethyl; hydroxyethoxy; lower alkoxy-(ethoxy)_n (n = 1 to 4); lower alkoxymethyl; cyanomethoxy; morpholine-4-yl; thiomorpholine-4-yl; 1-oxothiomorpholine-4-yl; 1,1-

- 15 dioxothiomorpholine-4-yl; 4-oxo-piperidine-1-yl; 4-alkoxy-piperidine-1-yl; 4-hydroxy-piperidine-1-yl; 4-hydroxyethoxy-piperidine-1-yl; 4-lower alkyl-piperazine-1-yl; alkoxycarbonyl; 2-dialkylamino-ethylsulfanyl; N,N-bis lower alkylamino lower alkyl; carbamoylmethyl; alkylsulfonyl; lower alkoxycarbonyl-lower alkyl; alkylcarboxy-lower alkyl; carboxy-lower alkyl; alkoxycarbonylmethylsulfanyl; carboxymethylsulfanyl; 1,4-
20 dioxo-8-aza-spiro[4.5]dec-8-yl; carboxy-lower alkoxy; cyano-lower alkyl; 2,3-dihydroxy-lower alkoxy; carbamoylmethoxy; 2-oxo-[1,3]-dioxolan-4-yl-lower alkoxy; (2-hydroxy-

- 2 -

lower alkyl)-lower alkyl amino; hydroxycarbamoyl-lower alkoxy; 2,2-dimethyl-tetrahydro-[1,3]dioxolo[4,5c]-pyrrol-5-yl; lower alkoxy-carbamoyl-lower alkoxy; 3R-hydroxypyrrolidin-1-yl; 3,4-dihydroxy-pyrrolidin-1-yl; 2-oxo-oxazolidin-3-yl; lower alkyl-carbamoylmethoxy; or aminocarbamoyl-lower alkoxy;

5 R^3 is a 5 or 6 membered aryl or heteroaryl which are optionally substituted by

halogen; cyano; nitro; azido; hydroxy; carboxy; morpholine-4-carbonyl; carbamoyl; thiocarbamoyl; N-hydroxycarbamoyl; trimethylsilyl-ethynyl;

or lower alkyl; lower alkoxy; halo-lower alkyl; 4-lower alkyl-piperazine-1-carbonyl; lower alkylcarbamoyl which are optionally substituted by amino, lower alkylamino, acylamino,
 10 oxo, hydroxy, lower alkoxy, lower alkylthio, or carboxy which is optionally esterified or amidated;

or an optionally substituted five-membered aromatic heterocycle, which may be optionally substituted by amino, lower alkylamino, acylamino, oxo, hydroxy, lower alkoxy, lower alkylthio, or carboxy which is optionally esterified or amidated, or lower alkyl which is
 15 optionally substituted by halogen, amino, lower alkylamino, acylamino, hydroxy, lower alkoxy, lower alkylthio, acyloxy, lower alkenoyl, lower alkylsulfinyl, lower alkylsulfonyl, cycloalkylsulfinyl, cycloalkylsulfonyl, hydroxyimino, alkoxyimino, carboxy which is optionally esterified or amidated, lower alkenyl, oxo, cyano, carbamoyloxy, sulfamoyl which is optionally substituted by lower alkyl, or amidino which is optionally substituted
 20 by lower alkyl, $-C(NRR')=NR''$ (where R, R' and R'' are hydrogen or lower alkyl) and their pharmaceutically acceptable addition salts.

It has surprisingly been found that the compounds of general formula I are metabotropic glutamate receptor antagonists. Compounds of formula I are distinguished by valuable therapeutic properties.

25 In the central nervous system (CNS) the transmission of stimuli takes place by the interaction of a neurotransmitter, which is sent out by a neuron, with a neuroreceptor.

L-glutamic acid, the most commonly occurring neurotransmitter in the CNS, plays a critical role in a large number of physiological processes. The glutamate-dependent stimulus receptors are divided into two main groups. The first main group forms ligand-controlled ion channels. The metabotropic glutamate receptors (mGluR) form the second
 30 main group and, furthermore, belong to the family of G-protein-coupled receptors.

At present, eight different members of these mGluR are known and of these some even have sub-types. On the basis of structural parameters, the different influences on the synthesis of secondary metabolites and the different affinity to low-molecular weight chemical compounds, these eight receptors can be sub-divided into three sub-groups:

5 mGluR1 and mGluR5 belong to group I, mGluR2 and mGluR3 belong to group II and mGluR4, mGluR6, mGluR7 and mGluR8 belong to group III.

Ligands of metabotropic glutamate receptors belonging to the group II can be used for the treatment or prevention of acute and/or chronic neurological disorders such as psychosis, schizophrenia, Alzheimer's disease, cognitive disorders and memory deficits.

10 Other treatable indications in this connection are restricted brain function caused by bypass operations or transplants, poor blood supply to the brain, spinal cord injuries, head injuries, hypoxia caused by pregnancy, cardiac arrest and hypoglycaemia. Further treatable indications are chronic and acute pain, Huntington's chorea, amyotrophic lateral sclerosis (ALS), dementia caused by AIDS, eye injuries, retinopathy, idiopathic parkinsonism or

15 parkinsonism caused by medicaments as well as conditions which lead to glutamate-deficiency functions, such as e.g. muscle spasms, convulsions, migraine, urinary incontinence, nicotine addiction, opiate addiction, anxiety, vomiting, dyskinesia and depressions.

Objects of the present invention are compounds of formula I and their pharmaceutically

20 acceptable salts per se and as pharmaceutically active substances, their manufacture, medicaments based on a compound in accordance with the invention and their production, as well as the use of the compounds in accordance with the invention in the control or prevention of illnesses of the aforementioned kind, and, respectively, for the production of corresponding medicaments.

25 Preferred compounds of formula I in the scope of the present invention are those in which R³ is phenyl substituted in meta position by cyano; halogen; or imidazolyl which is optionally substituted by lower alkyl or methylsulfanyl; 1,2,3-triazolyl; 1,2,4-triazolyl; or isoxazolyl which is optionally substituted by lower alkyl.

The following are examples of such compounds:

30 3-(8-Chloro-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile;

3-[8-(4-Methyl-piperazin-1-yl)-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile;

- 3-(8-Chloro-4-oxo-7-phenyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile;
[4-(3-Cyano-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-ylsulfanyl]-acetic acid methyl ester;
2-[4-(3-Cyano-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-yl]-acetamide;
3-(8-Methoxy-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile;
3-(8-Cyanomethyl-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile;
4-(3-Iodo-phenyl)-7-(2-methoxy-ethoxy)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
4-(3-Imidazol-1-yl-phenyl)-7-(2-methoxy-ethoxy)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
[RS]-3-[4-Oxo-8-(2-oxo-[1,3]dioxolan-4-ylmethoxy)-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile;
7-Hydroxymethyl-4-(3-imidazol-1-yl-phenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
[4-(3-Imidazol-1-yl-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-yloxy]-acetonitrile;
8-(4-Fluoro-phenylethynyl)-7-hydroxymethyl-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
7-(2-Hydroxy-ethoxy)-4-(3-imidazol-1-yl-phenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
8-(4-Fluoro-phenyl)-7-[4-(2-hydroxy-ethoxy)-piperidin-1-yl]-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
8-(4-Fluoro-phenyl)-7-hydroxy-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
8-(2-Fluoro-phenyl)-7-methoxy-4-[3-(2-methyl-imidazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
8-(2-Fluoro-phenyl)-7-hydroxy-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
8-(2-Fluoro-phenyl)-7-hydroxy-4-[3-(2-methyl-imidazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
8-(2-Fluoro-phenyl)-7-hydroxy-4-[3-(2-methylsulfanyl-imidazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
8-(2,5-Difluoro-phenyl)-7-methoxy-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one;

- 8-(2-Fluoro-phenyl)-7-hydroxy-4-[3-(3-methyl-isoxazol-5-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
- 3-[7-(2,5-Difluoro-phenyl)-8-hydroxy-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile;
- 5 8-(4-Fluoro-phenylethynyl)-7-hydroxy-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one; and
- 8-(4-Fluoro-phenylethynyl)-7-hydroxy-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one.

Compounds of formula I, wherein R³ is thiophenyl, preferably thiophen-2-yl, which is
10 optionally substituted by cyano or halogen; or R³ is pyridinyl, preferably pyridin-4-yl, which is optionally substituted in 2-position by cyano or halogen, or wherein R³ is thiazolyl which is optionally substituted in 2-position with imidazolyl or 4-methylimidazolyl, are also preferred.

The following compounds are particularly preferred:

- 15 5-[7-(2-Fluoro-phenyl)-8-methoxy-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-thiophene-2-carbonitrile;
- 2-[7-(2-Fluoro-phenyl)-8-hydroxy-4-oxo-4,5-dihydro-1H-benzo[b][1,4]diazepin-2-yl]-thiophene-3-carbonitrile;
- 4-[7-(2-Fluoro-phenyl)-8-methoxy-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-
20 pyridine-2-carbonitrile;
- 4-[7-(4-Fluoro-phenyl)-8-hydroxy-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-pyridine-2-carbonitrile;
- 4-[7-(2-Fluoro-phenyl)-8-hydroxy-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-pyridine-2-carbonitrile; and
- 25 8-(2-Fluoro-phenyl)-4-[2-(4-methyl-imidazol-1-yl)-thiazol-4-yl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one.

All tautomeric forms of the compounds of the invention are also embraced herewith.

The term "lower alkyl" used in the present description denotes straight-chain or branched saturated hydrocarbon residues with 1-7 carbon atoms, preferably with 1-4 carbon atoms,
30 such as methyl, ethyl, n-propyl, i-propyl and the like.

The term "lower cycloalkyl" used in the present description denotes cyclic saturated hydrocarbon residues with 3-5 carbon atoms, preferably with 3 carbon atoms, such as cyclopropyl.

The term "lower alkoxy" denotes a lower alkyl residue in the sense of the foregoing definition bonded via an oxygen atom.


The term "halogen" embraces fluorine, chlorine, bromine and iodine.

The term "5 or 6 membered aryl or heteroaryl" embraces phenyl, thiophenyl, pyridine,
5 partially hydrated pyridine.

The expression "five-membered aromatic heterocycle" embraces, furan, thiazol, imidazol, pyrazol, 1,3-thiazol, 1,3-oxazol, 1,2-oxazol, 1,2-thiazol, 1,2,3-triazol, 1,2,4-triazol, 1,2,4-oxadiazol, 1,2,3-oxadiazol, 1,2,4-thiadiazol, 1,2,3-thiadiazol and tetrazol.

The compounds of general formula I and their pharmaceutically acceptable salts can be
10 manufactured according to the following methods:

Scheme A



According to scheme A, compounds of general formula I, in which X , R^1 , R^2 and R^3 are as described above, can be prepared from compounds of general formula II via an acylation-deprotection-cyclization sequence:
15

For example reacting compounds of general formula II with a dioxinone IV, in which R^3 is as described above, in an inert solvent such as toluene or xylene at elevated temperatures, preferably between 80 °C and 160 °C gives rise to compounds of general formula III.


Alternatively, compounds of general formula III can also be prepared by for example
20 reaction of a compound of general formula II with a β -ketoester (general formula IVa), in which R^3 is as described above using the same conditions as described for the reaction with the dioxinones.

Afterwards, cleaving the BOC protecting group in compounds of general formula III and concomitant cyclization of the deprotected compound yields the desired compounds of general formula I. Any other suitable amino protecting group, such as e.g. Fmoc or benzyloxycarbonyl (Z), can be alternatively used instead of the BOC group.

- 5 The deprotection-cyclization step can be carried out by treating the compounds of general formula III with for example a Bronsted acid such as trifluoroacetic acid in an inert solvent such as dichloromethane (DCM). The reaction is preferably carried out at temperatures between 0 °C and 50 °C. It may be advantageous to use also anisole or 1,3-dimethoxybenzene as a carbocation scavenger in the reaction mixture.

10

Scheme B




According to scheme B, compounds of general formula II in which R^1 is as described above for compounds where X is a single bond and R^2 is as described above, can be prepared by different routes depending on the nature of R^1 from the iodo-compounds of general formula V, in which R^2 is as described above. As shown in scheme B, the key steps are

5 coupling reactions of Suzuki- and Stille-type in presence or absence of carbonmonoxide. The exact conditions for the respective compounds of general formula II can be found in the experimental part.

Compounds of general formula V, in which R^2 is as described above, can be prepared by different routes depending on the individual residue R^2 :

10

Scheme C

As shown in scheme C, compounds of general formula Va, in which R^2 is lower alkyl, halogen or alkoxy-carbonyl, can be prepared from the known compounds of general formula XI by iodination and subsequent protection of the synthetic intermediates with

15 general formula XII.

The iodination step can be carried out by for example using iodine monochloride in acetic acid in the presence of sodium acetate. The reaction can be for example carried out at temperatures between 20 °C and 80 °C

The protection of the amino function can be achieved by for example reacting compounds


20 of general formula XII with di-tert.-butyl-carbonate in the presence of a base such as cesium carbonate. The reaction can be carried out in polar solvents such as acetone or butanone and the like at temperatures between 20 °C and 60 °C.

As shown in scheme D, compounds of general formula Vb and Vc, in which R^2 is attached via a sulfur- or nitrogen-atom (R^2 represents for example morpholin-4-yl;


25 thiomorpholino-4-yl; dialkylamino; carboxymethylsulfanyl etc), respectively, can be prepared from the intermediate XIII by a nucleophilic substitution reaction with the respective amines or mercaptanes in the presence of a suitable base.

Scheme D

Nitrogen nucleophiles



Sulfur nucleophiles




The reaction is preferably carried out in a polar, aprotic solvent such as dimethyl formamide, N-methyl-pyrrolidone or dimethyl sulfoxide and the like. The base can be
 5 selected from the sterically hindered amines such as Hünig's base, alkoxides such as sodium methoxide and tert.-butoxide, or hydrides such as sodium hydride. The reaction can be performed at temperatures between 20 °C and 110 °C, depending on the individual compounds to be synthesized.

Compounds of general formula Vd in which R² is attached via an oxygen atom (R²
 10 represents for example lower alkoxy, lower halo-alkoxy, lower cyclo-alkoxy, lower alkoxy-lower alkoxy; etc.) can be prepared as for example shown in scheme E:

Scheme E


Oxygen nucleophiles



GP B, method a: diphosgene, EtOAc, 77 °C; then t-BuOH

GP B, method b: Boc_2O , Cs_2CO_3 , 2-butanone, 52 °CGP B, method c: i) Boc_2O , DMAP, THF; ii) TFA, DCM, 0 °C

Oxygen alkylation



by a nucleophilic aromatic substitution reaction with the respective alcohol in the presence of a suitable base and subsequent protection of the amino function. The base can be selected from the class of Bronsted bases such as potassium hydroxide and the like. The reaction is preferably carried out in a polar, aprotic solvent such as dimethyl formamide, N-methyl-pyrrolidone or dimethyl sulfoxide and the like at temperatures between 20 °C and 100 °C.

The protection of the amino function can be achieved by for example reacting compounds of general formula XV with di-tert.-butoxy carbonate in the presence of a base such as cesium carbonate. The reaction can be carried out in polar solvents such as acetone or butanone and the like at temperatures between 20 °C and 60 °C.

Another method to achieve this protection step is to transform first the amino function in a compound with general formula XV into an isocyanate by reaction with phosgene or a phosgene equivalent in the presence of a suitable base, which is then treated with tert.-butyl-alcohol to give the desired compounds of general formula Vd.

Another suitable method to achieve this protection step is to transform first the amino function in a compound with general formula XV into the corresponding di-Boc compound by reaction with excess di-tert.-butoxy carbonate in the presence of 4-dimethylaminopyridine (DMAP), which is then treated with 2 eq. TFA in dichloromethane to give the desired compounds of general formula Vd.


This reversal of steps, i.e. performing first the nucleophilic aromatic substitution on the key intermediate XIV and second protection of the amino-function as shown in synthetic scheme E can also be applied to those compounds with the general formula Vb and Vc (synthetic scheme D).

- 5 Yet another method of preparing compounds of the general formula Vd is using the O-allyl compound XIX and performing a deallylation-alkylation sequence as outlined in scheme E. The deallylation is preferably carried out by transition-metal catalyzed isomerisation, e.g. in the presence of Rhodium(I)-salts like for example Wilkinson's catalyst $[(PPh_3)_3RhCl]$ or Palladium(II)-salts such as $[(PPh_3)_2PdCl_2]$, followed by aqueous acid hydrolysis of the
- 10 resulting vinyl ether. An example for this procedure can be found in *J. Org. Chem.* 1973, 38, 3224. The alkylation of the phenol XX to the desired compound of the general formula Vd can be carried out with electrophilic reagents of the general formula R-X, in which R has the meaning of lower alkyl, lower alkenyl, alkyl acetate or benzyl and X represents a leaving group, for example iodide, bromide, methanesulfonate or tolylsulfonate, in a
- 15 suitable solvent in the presence of a base. The reaction is preferably carried out in polar, aprotic solvents, for example chlorinated solvents such as dichloromethane, chloroform or dichloroethane, or amides, for example dimethylformamide, dimethylacetamide and N-methyl-pyrrolidone, or sulfoxides, for example dimethyl sulfoxide. The base can be selected from the sterically hindered amines such as Hünig's base, alkoxides such as
- 20 sodium methoxide and tert.-butoxide, hydrides such as sodium hydride, hydroxides such as potassium hydroxide, carbonates such as potassium carbonate or hydrogen carbonates such as potassium hydrogen carbonate. The reaction can be performed at temperatures between -20 °C and 80 °C, depending on the individual compounds to be synthesized. For the synthesis of an O-tert.-butyl compound with the general formula Vd the phenol XX
- 25 can be treated with DMF-di-tert.-butylacetal in toluene or benzene at 80 °C as described in *Synthesis* 1983, 135.


According to synthetic scheme F,

Scheme F

Carbon nucleophiles



Cl-moiety




- 5 compounds of general formula Ve and Vf in which R^2 is attached via a carbon atom (R^2 represents for example lower alkoxy-carbonyl-methyl; cyano-methyl, etc.) can be prepared from compound XIII or XIV by for example reaction with a malonic acid ester or -half-ester in the presence of a base followed by the removal of one of the alkyl carboxylates via decarboxylation. The exact reaction conditions vary with the identity of the individual
- 10 compounds and are described in the examples.

The key intermediates XIII and XIV can be prepared as already described in scheme C.

- For the one-carbon-moiety bearing compounds of the general formula Vh to VI, the synthesis starts from known methyl 3-amino-4-nitrobenzoate. Standard iodination as described in synthetic scheme C leads to the iodide XXI, which in turn can be protected with the Boc-group. The reduction of the methyl ester can for example be performed by
- 15 treatment with lithium borohydride, sodium borohydride or diisobutylaluminumhydride in an aprotic solvent like for example THF, ether or toluene. The presence of an alcohol such as methanol, ethanol or isopropanol can be advantageous. The reduction is preferably carried out at temperatures between -20°C and 0°C . Further functionalization, like for
- 20 example conversion into a chloride (Vk), of the resulting benzylic alcohol Vh follows

standard procedures known to someone skilled in the art. The exact reaction conditions vary with the identity of the individual compounds and are described in the examples.

Scheme G




5 According to scheme G, compounds of general formula II in which R¹ is as described above for compounds where X is an ethynediyl group can be prepared by different routes from the iodo-compounds V, depending on the nature of R¹ and R². As shown in scheme G, the transformation can for example be carried out

a) by directly attaching the R¹-alkynediyl-substituent to a compound of general formula V
 10 via a Sonogashira-type coupling followed by the reduction of the nitro group or

b) by two stepwise Sonogashira-type couplings, in which first trimethylsilyl-acetylene is coupled to a compound of general formula V to yield, after deprotection with sodium hydroxide in methanol, the intermediate XVIII which then can be transformed via a second Sonogashira-type coupling with the appropriate reactant R¹-I, R¹-Br or R¹-OSO₂CF₃ and reduction of the nitro group to the desired compounds of general formula II.
 15


The exact conditions for the respective compounds can be found in the experimental part.

Scheme H



- 5 According to Scheme H, the dioxinones and β -keto esters building blocks with the general formula IV and IVa can be prepared by methods known to someone skilled in the art from the corresponding carboxylic acid derivatives R^3-COR , i.e. free acids, methyl or ethyl esters and acid chlorides. The exact conditions for the corresponding compounds can be found in the experimental part.
- 10 Another synthetic route to prepare compounds of general formula I in which R^1, R^2 and X have the meaning as described above and R^3 is a carbamide of general formula $C(O)NR^4R^5$, in which R^4 and R^5 is hydrogen, lower alkyl or R^4 and R^5 together form a morpholino-residue or a N-methyl-piperazine, is outlined in scheme I:

Scheme I



The exact conditions for the respective compounds can be found in the experimental part.

The pharmaceutically acceptable salts can be manufactured readily according to methods known per se and taking into consideration the nature of the compound to be converted into a salt. Inorganic or organic acids such as, for example, hydrochloric acid, hydrobromic acid, sulphuric acid, nitric acid, phosphoric acid or citric acid, formic acid, fumaric acid, maleic acid, acetic acid, succinic acid, tartaric acid, methanesulphonic acid, p-toluenesulphonic acid and the like are suitable for the formation of pharmaceutically acceptable salts of basic compounds of formula I. The compounds of formula I and their pharmaceutically acceptable salts are metabotropic glutamate receptor antagonists and can be used for the treatment or prevention of acute and/or chronic neurological disorders, such as psychosis, schizophrenia, Alzheimer's disease, cognitive disorders and memory deficits. Other treatable indications are restricted brain function caused by bypass operations or transplants, poor blood supply to the brain, spinal cord injuries, head injuries, hypoxia caused by pregnancy, cardiac arrest and hypoglycaemia. Further treatable indications are acute and chronic pain, Huntington's chorea, ALS, dementia caused by AIDS, eye injuries, retinopathy, idiopathic parkinsonism or parkinsonism caused by medicaments as well as conditions which lead to glutamate-deficient functions, such as e.g. muscle spasms, convulsions, migraine, urinary incontinence, nicotine addiction, psychoses, opiate addiction, anxiety, vomiting, dyskinesia and depression.

The compounds of formula I and pharmaceutically acceptable salts thereof can be used as medicaments, e.g. in the form of pharmaceutical preparations. The pharmaceutical preparations can be administered orally, e.g. in the form of tablets, coated tablets, dragées, hard and soft gelatine capsules, solutions, emulsions or suspensions. However, the administration can also be effected rectally, e.g. in the form of suppositories, or parenterally, e.g. in the form of injection solutions.

The compounds of formula I and pharmaceutically acceptable salts thereof can be processed with pharmaceutically inert, inorganic or organic carriers for the production of pharmaceutical preparations. Lactose, corn starch or derivatives thereof, talc, stearic acid or its salts and the like can be used, for example, as such carriers for tablets, coated tablets, dragées and hard gelatine capsules. Suitable carriers for soft gelatine capsules are, for example, vegetable oils, waxes, fats, semi-solid and liquid polyols and the like; depending on the nature of the active substance no carriers are, however, usually required in the case of soft gelatine capsules. Suitable carriers for the production of solutions and syrups are, for example, water, polyols, sucrose, invert sugar, glucose and the like. Adjuvants, such as alcohols, polyols, glycerol, vegetable oils and the like, can be used for aqueous injection

- 16 -

solutions of water-soluble salts of compounds of formula I, but as a rule are not necessary. Suitable carriers for suppositories are, for example, natural or hardened oils, waxes, fats, semi-liquid or liquid polyols and the like.

In addition, the pharmaceutical preparations can contain preservatives, solubilizers, stabilizers, wetting agents, emulsifiers, sweeteners, colorants, flavorants, salts for varying the osmotic pressure, buffers, masking agents or antioxidants. They can also contain still other therapeutically valuable substances.

As mentioned earlier, medicaments containing a compound of formula I or a pharmaceutically acceptable salt thereof and a therapeutically inert excipient are also an object of the present invention, as is a process for the production of such medicaments which comprises bringing one or more compounds of formula I or pharmaceutically acceptable salts thereof and, if desired, one or more other therapeutically valuable substances into a galenical dosage form together with one or more therapeutically inert carriers.

The dosage can vary within wide limits and will, of course, be fitted to the individual requirements in each particular case. In general, the effective dosage for oral or parenteral administration is between 0.01-20 mg/kg/day, with a dosage of 0.1-10 mg/kg/day being preferred for all of the indications described. The daily dosage for an adult human being weighing 70 kg accordingly lies between 0.7-1400 mg per day, preferably between 7 and 700 mg per day.

The present invention relates also to the use of compounds of formula I and of pharmaceutically acceptable salts thereof for the production of medicaments, especially for the control or prevention of acute and/or chronic neurological disorders of the aforementioned kind.

The compounds of the present invention are group II mGlu receptor antagonists. The compounds show activities, as measured in the assay described below, of 50 μ M or less, typically 3 μ M or less, and ideally of 0.5 μ M or less. In the table below are described some specific pKi values of preferred compounds.

Compound	K _i mGlu2 (μM)
3-(8-Chloro-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile	0.028
3-[8-(4-Methyl-piperazin-1-yl)-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile	0.305
3-(8-Chloro-4-oxo-7-phenyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile	0.120
[4-(3-Cyano-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-ylsulfanyl]-acetic acid methyl ester	0.051
2-[4-(3-Cyano-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-yl]-acetamide	0.037
3-(8-Methoxy-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile	0.046
3-(8-Cyanomethyl-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile	0.016
4-(3-Iodo-phenyl)-7-(2-methoxy-ethoxy)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.021
4-(3-Imidazol-1-yl-phenyl)-7-(2-methoxy-ethoxy)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.012
[RS]-3-[4-Oxo-8-(2-oxo-[1,3]dioxolan-4-ylmethoxy)-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile	0.035
7-Hydroxymethyl-4-(3-imidazol-1-yl-phenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.018
[4-(3-Imidazol-1-yl-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-yloxy]-acetonitrile	0.009

8-(4-Fluoro-phenylethynyl)-7-hydroxymethyl-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.009
7-(2-Hydroxy-ethoxy)-4-(3-imidazol-1-yl-phenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.032
8-(4-Fluoro-phenyl)-7-[4-(2-hydroxy-ethoxy)-piperidin-1-yl]-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.100
8-(4-Fluoro-phenyl)-7-hydroxy-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.007
8-(2-Fluoro-phenyl)-7-methoxy-4-[3-(2-methyl-imidazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.138
5-[7-(2-Fluoro-phenyl)-8-methoxy-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-thiophene-2-carbonitrile	0.168
4-[7-(2-Fluoro-phenyl)-8-methoxy-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-pyridine-2-carbonitrile	0.033
8-(2-Fluoro-phenyl)-7-hydroxy-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.028
8-(2-Fluoro-phenyl)-7-hydroxy-4-[3-(2-methyl-imidazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.108
8-(2-Fluoro-phenyl)-7-hydroxy-4-[3-(2-methylsulfanyl-imidazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.021
8-(2,5-Difluoro-phenyl)-7-methoxy-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.012
8-(2-Fluoro-phenyl)-7-hydroxy-4-[3-(3-methyl-isoxazol-5-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.015
3-[7-(2,5-Difluoro-phenyl)-8-hydroxy-4-oxo-4,5-dihydro-	0.006

3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile	
8-(4-Fluoro-phenylethynyl)-7-hydroxy-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.013
8-(4-Fluoro-phenylethynyl)-7-hydroxy-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one	0.010

[³H]-LY354740 binding on mGlu2 transfected CHO cell membranes

Transfection and cell culture

cDNA encoding the rat mGlu2 receptor protein in pBluescript II was obtained from Prof. S. Nakanishi (Kyoto, Japan), and subcloned into the eukaryotic expression vector pcDNA I-amp from Invitrogen (NV Leek, The Netherlands). This vector construct (pcD1mGR2) was co-transfected with a psvNeo plasmid encoding the gene for neomycin resistance, into CHO cells by a modified calcium phosphate method described by Chen & Okayama (1988). The cells were maintained in Dulbecco's Modified Eagle medium with reduced L-glutamine (2 mM final concentration) and 10% dialysed foetal calf serum from Gibco BRL (Basel, Switzerland). Selection was made in the presence of G-418 (1000 ug/ml final). Clones were identified by reverse transcription of 5 µg total RNA, followed by PCR using mGlu2 receptor specific primers 5'-atcactgcttggttctggcactg-3' and 5'-agcatcactgtgggtggcataggagc-3' in 60 mM Tris HCl (pH 10), 15 mM (NH₄)₂SO₄, 2 mM MgCl₂, 25 units/ml Taq Polymerase with 30 cycles annealing at 60 °C for 1 min., extension at 72 °C for 30 s, and 1 min. 95 °C denaturation.

Membrane preparation

Cells, cultured as above, were harvested and washed three times with cold PBS and frozen at -80 °C. The pellet was resuspended in cold 20 mM HEPES-NaOH buffer containing 10 mM EDTA (pH 7.4), and homogenised with a polytron (Kinematica, AG, Littau, Switzerland) for 10s at 10 000rpm. After centrifugation for 30 min. at 4 °C, the pellet was washed once with the same buffer, and once with cold 20 mM HEPES-NaOH buffer containing 0.1 mM EDTA, (pH 7.4). Protein content was measured using the Pierce method (Socochim, Lausanne, Switzerland) using bovine serum albumin as standard.

[³H]-LY354740 binding

After thawing, the membranes were resuspended in cold 50mM Tris-HCl buffer containing 2 mM MgCl₂ and 2 mM CaCl₂, (pH 7) (binding buffer). The final concentration of the

membranes in the assays was 25 µg protein/ml. Inhibition experiments were performed with membranes incubated with 10 nM [³H]-LY354740 at room temperature, for 1 hour, in presence of various concentrations of the compound to be tested. Following the incubations, membranes were filtered onto Whatmann GF/C glass fiber filters and washed 5 times with cold binding buffer. Non specific binding was measured in the presence of 10 µM DCG IV. After transfer of the filters into plastic vials containing 10 ml of Ultima-gold scintillation fluid (Packard, Zürich, Switzerland), the radioactivity was measured by liquid scintillation in a Tri-Carb 2500 TR counter (Packard, Zürich, Switzerland).

Data analysis.

- 10 The inhibition curves were fitted with a four parameter logistic equation giving IC₅₀ values, and Hill coefficients.

EXAMPLES

The following examples relate to the preparation of the (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl esters (Synthetic Scheme C):

15 General procedure A

Preparation of 4-iodo-2-nitroanilines by iodination of 2-nitroanilines [according to Wilson, J. Gerald; Hunt, Frederick C. *Aust. J. Chem.* 1983, 36, 2317-25]

- To a stirred solution of the 2-nitroaniline (1.0 mol) in HOAc (500 mL) containing anhydrous NaOAc (93-103 g, 1.125-1.25 mol), iodine monochloride (59-66 mL, 1.125-1.25 mol) in HOAc (300 mL) was added over 60 min. The reaction mixture was heated to the given temperature until thin layer chromatography (tlc) indicated complete conversion of the starting material, stirred for another 30 min at 23 °C, then diluted slowly with H₂O (1000 mL) which caused the separation of the crystalline product. Stirring was continued for 1 h and the product was filtered off, washed free of HOAc and dried in vacuum at 60 °C.

Example A1

5-Chloro-4-iodo-2-nitro-phenylamine

Prepared from 5-chloro-2-nitroaniline by iodination with iodine monochloride in HOAc/NaOAc according to the general procedure A (80 °C). Obtained as an orange solid.

- 30 MS (EI) 298 (M⁺) and 300 [(M+2)⁺]; mp 202-203 °C (dec.).

Example A2

4-Iodo-5-methyl-2-nitro-phenylamine

Prepared from 5-methyl-2-nitroaniline by iodination with iodine monochloride in HOAc/NaOAc according to the general procedure A (80 °C). Obtained as a red solid.

5 MS (EI) 278 (M⁺); mp 154 °C (dec.).

Example A3

5-Amino-2-iodo-4-nitrobenzoic acid methyl ester

Prepared from 3-amino-4-nitrobenzoic acid methyl ester (22.25 g, {CAS-No. [99512-09-1]}; prepared in two steps as follows: 3-hydroxy-4-nitrobenzoic acid (30 g, 164 mmol),
10 NH₄Cl (21.91 g, 410 mmol) in 25% aq. NH₃ (180 mL) was heated in a steel autoclave at 160 °C for 7 h (internal pressure: 23 bar). Cooled to 23 °C and evaporated to dryness. Taken up in H₂O (200 mL), adjusted pH with conc. H₂SO₄ to pH 1, saturated with NaCl and extracted with EtOAc (6 x 750 mL), dried combined organic layer over MgSO₄. Filtration and removal of the solvent in vacuum left the sufficiently pure 3-amino-4-
15 nitrobenzoic acid (22.26 g, 75 %) as an orange solid. This material was suspended in MeOH (500 mL), conc. H₂SO₄ (3 mL) was added and the mixture was heated to 65 °C for 2.5 days. The solvent was removed in vacuum, the solid residue taken up in EtOAc, washed with sat. NaHCO₃-sol. and brine, followed by drying over MgSO₄. Removal of the solvent left the sufficiently pure 3-amino-4-nitrobenzoic acid methyl ester (22.25 g, 93 %) as an
20 orange solid.) by iodination with iodine monochloride in HOAc/NaOAc according to the general procedure A (35 °C). Obtained as an orange solid (29.38 g, 80 %).

MS (EI) 322 (M⁺); mp 168 °C (dec.).

General procedure B:

Preparation of (2-nitro-phenyl)-carbamic acid tert.-butyl esters from 2-nitroanilines

25 Method a: To a solution of diphosgene (4.1 mL, 34.1 mmol) in EtOAc (40 mL) at 0 °C was added a solution of the 4-iodo-2-nitroaniline (45.5 mmol) in EtOAc (200-500 mL), and the mixture was heated to reflux for 18 h. The solvent was removed in vacuum to leave a brown solid, which was triturated with hot hexane (200 mL). The solid material was filtered off and the filtrate was concentrated under reduced pressure to leave the pure 4-
30 iodo-2-nitrophenylisocyanate as a yellow solid. This material was refluxed in a mixture of excess tert.-BuOH in CH₂Cl₂ for 2.5 h. Removal of the solvent left an orange solid which

was purified by silica gel column chromatography with hexane/EtOAc to give the (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester as a yellow solid.

Method b: To a mixture of the 4-iodo-2-nitroaniline (142 mmol) and cesium carbonate (55.5 g, 170 mmol) in 2-butanone (740 mL) was dropwise added a solution of Boc_2O (37.8 g, 173 mmol) in 2-butanone (170 mL) and the resulting mixture was stirred at 52 °C for 26 h. The solvent was removed in vacuum, the residue was treated with a mixture of H_2O (240 mL) and MeOH (240 mL) and extracted with hexane (3 x 500 mL). The combined hexane layer was washed with brine (200 mL) and all aqueous layers were reextracted with hexane (300 mL). All combined hexane layers were dried over MgSO_4 , filtered and the solvent was removed in vacuum to give an orange solid, which was purified by silica gel column chromatography with hexane/EtOAc to give the (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester as a yellow solid.

Method c: To a solution of the 4-iodo-2-nitroaniline (550 mmol) and DMAP (1.22 g, 10 mmol) in THF (1000 mL) at 23 °C was dropwise added within 70 min a solution of Boc_2O (246 g, 1128 mmol) in THF (500 mL) and stirring was continued at 23 °C for 75 min. The entire mixture was evaporated to dryness and dried at HV to leave a dark brown solid (253.59 g). This material was dissolved in DCM (1100 mL), cooled to 0 °C and TFA (84 mL, 1100 mmol) was added dropwise. The mixture was stirred at 0 °C for 2 h, poured into icecold sat. NaHCO_3 -sol., extracted with DCM, washed with brine and dried over MgSO_4 . Removal of the solvent in vacuum left a dark brown solid (199.71 g) which was coated on silica gel and purified by silica gel column chromatography with hexane/EtOAc to give the (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester as a yellow solid.

Example B1

(5-Chloro-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester

Prepared the isocyanate from 5-chloro-4-iodo-2-nitro-phenylamine (Example A1) (7.0 g, 23.45 mmol) with diphosgene (2.12 mL, 17.6 mmol) in EtOAc (30 mL), followed by treatment with tert.-BuOH (100 mL) in CH_2Cl_2 (100 mL) according to the general procedure B (method a). Obtained as a yellow solid (7.1 g, 76 %).

MS (EI) 398 (M^+) and 400 [$(\text{M}+2)^+$]; mp 82-84 °C.

Example B2

(4-Iodo-5-methyl-2-nitro-phenyl)-carbamic acid tert.-butyl ester

Prepared the isocyanate from 4-iodo-5-methyl-2-nitro-phenylamine (Example A2) (13.51 g, 48.6 mmol) with diphosgene (4.4 mL, 36.4 mmol) in EtOAc (50 mL), followed by treatment with tert.-BuOH (150 mL) and CH₂Cl₂ (150 mL) according to the general procedure B (method a). Obtained as a yellow solid (14.1 g, 77 %).

5 MS (EI) 378 (M⁺); mp 99-100 °C.

Example B3

5-tert.-Butoxycarbonylamino-2-iodo-4-nitro-benzoic acid methyl ester

Prepared the isocyanate from 5-amino-2-iodo-4-nitro-benzoic acid methyl ester (Example A3) (5.5 g, 17 mmol) with diphosgene (1.55 mL, 13 mmol) in EtOAc (135 mL), followed
10 by treatment with tert.-BuOH (20 mL) and CH₂Cl₂ (70 mL) according to the general procedure B (method a). Obtained as a yellow solid (5.2 g, 72 %).

MS (ISP) 440 [(M+NH₄)⁺]; mp 126 °C.

Example B4

(5-Allyloxy-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester

15 Prepared the isocyanate from 5-allyloxy-4-iodo-2-nitro-phenylamine (Example E1) (9.0 g, 28.2 mmol) with diphosgene (2.6 mL, 21.2 mmol) in EtOAc (150 mL), followed by treatment with tert.-BuOH (80 mL) and CH₂Cl₂ (80 mL) according to the general procedure B (method a). Obtained as a yellow solid (9.16 g, 77 %).

MS (ISP) 421 [(M+H)⁺] and 438 [(M+NH₄)⁺]; mp 93-95 °C.

20 Example B5

(4-Iodo-5-methoxy-2-nitro-phenyl)-carbamic acid tert.-butyl ester

Prepared the isocyanate from 4-iodo-5-methoxy-2-nitro-phenylamine (Example E2) (2.85 g, 9.69 mmol) with diphosgene (0.88 mL, 7.27 mmol) in EtOAc (52 mL), followed by treatment with tert.-BuOH (25 mL) and CH₂Cl₂ (25 mL) according to the general
25 procedure B (method a). Obtained as a yellow solid (3.0 g, 79 %).

MS (EI) 394 (M⁺); mp 171 °C.

Example B6

[4-Iodo-5-(2-methoxy-ethoxy)-2-nitro-phenyl]-carbamic acid tert.-butyl ester

Prepared the isocyanate from 4-iodo-5-(2-methoxy-ethoxy)-2-nitro-phenylamine (Example E3) (2.73 g, 8.08 mmol) with diphosgene (0.8 mL, 6.06 mmol) in EtOAc (50 mL), followed by treatment with tert.-BuOH (25 mL) and CH₂Cl₂ (25 mL) according to the general procedure B (method a). Obtained as a yellow solid (3.0 g, 86 %).

MS (ISP) 439 [(M+H)⁺], 456 [(M+NH₄)⁺] and 461 [(M+Na)⁺]; mp 109-111 °C.

Example B7

[4-Iodo-5-(2-[2-[2-(2-methoxy-ethoxy)-ethoxy]-ethoxy]-ethoxy)-2-nitro-phenyl]-carbamic acid tert.-butyl ester

Prepared the isocyanate from 4-Iodo-5-(2-[2-[2-(2-methoxy-ethoxy)-ethoxy]-ethoxy]-ethoxy)-2-nitro-phenylamine (Example E4) (8.0 g, 17 mmol) with diphosgene (1.54 mL, 13 mmol) in EtOAc (100 mL), followed by treatment with tert.-BuOH (25 mL) and CH₂Cl₂ (25 mL) according to the general procedure B (method a). Obtained as a yellow oil (8.6 g, 89 %).

MS (ISP) 588 [(M+NH₄)⁺].

Example B8

(RS)-[5-(2,2-Dimethyl-[1,3]dioxolan-4-ylmethoxy)-2-nitro-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester

Prepared from (RS)-5-(2,2-dimethyl-[1,3]dioxolan-4-ylmethoxy)-2-nitro-4-phenylethynyl-phenylamine (Example K14) (2.678 g, 5.7 mmol), Cs₂CO₃ (2.23 g, 6.8 mmol) and Boc₂O (1.52 g, 7.0 mmol) in 2-butanone (36.5 mL) at 52 °C according to the general procedure B (method b). Obtained as a yellow foam (2.0 g, 75 %).

MS (ISP) 469 [(M+H)⁺] and 486 [(M+NH₄)⁺]; mp 32 °C.

Example B9 [Ve (R=CN; R''=Boc)]

(5-Cyanomethyl-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester

Prepared the isocyanate from (5-amino-2-iodo-4-nitro-phenyl)-acetonitrile (Example Vf (R=CN; R''=H)) (5.15 g, 17 mmol) with diphosgene (2.05 mL, 17 mmol) in EtOAc (150

mL), followed by treatment with tert.-BuOH (25 mL) and CH₂Cl₂ (25 mL) according to the general procedure B (method a). Obtained as a yellow solid (4.0 g, 58 %).

MS (ISN) 402 [(M-H)⁻]; mp 124-126 °C.

Example B10

5 [5-(2-tert.-Butoxy-ethoxy)-4-iodo-2-nitro-phenyl]-carbamic acid tert.-butyl ester

Prepared the di-Boc-compound from 5-(2-tert.-butoxy-ethoxy)-4-iodo-2-nitro-phenylamine (Example E6) (13.9 g, 36.6 mmol) and Boc₂O (16.35 g, 75 mmol), followed by treatment with 2 eq. TFA in CH₂Cl₂ according to the general procedure B (method c). Obtained as a yellow solid (14.96 g, 85 %).

10 MS (ISP) 481 [(M+H)⁺]; mp 113-116 °C.

Example B11

[4-Iodo-5-(4-methoxy-benzyloxy)-2-nitro-phenyl]-carbamic acid tert.-butyl ester

Prepared the di-Boc-compound from 4-iodo-5-(4-methoxy-benzyloxy)-2-nitro-phenylamine (Example E7) (2.88 g, 7.20 mmol) and Boc₂O (3.30 g, 15.12 mmol), followed
15 by treatment with 2 eq. TFA in CH₂Cl₂ according to the general procedure B (method c). Obtained as a waxy yellow solid (1.74 g).

MS (ISN) 499 [(M-H)⁻].

The following procedures relate to the preparation of those (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl esters bearing nitrogen substituents in the 5-position (Scheme D).

20 General procedure C:

Preparation of 5-N-substituted-(4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl esters:

Method a: from (5-chloro-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester

(5-Chloro-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B1) was stirred with the desired amine optionally with toluene or DMSO and/or DIPEA at temperature

25 from 23 °C to 100-130 °C until tlc indicated complete disappearance of the chloride. The reaction was cooled to 23 °C poured into ice-water, the precipitate was filtered off, washed with water and dried in vacuum. In cases where the product did not precipitate, the mixture was extracted with EtOAc, washed with water and brine, dried over Na₂SO₄. Filtration and removal of the solvent in vacuum left a crude product, which was purified by silica gel

30 column chromatography with hexane/EtOAc to give the pure title compound.

Method b: from 5-chloro-4-iodo-2-nitro-phenylamine

A mixture of 5-chloro-4-iodo-2-nitro-phenylamine (Example A1) (1.49 g, 5.0 mmol), the desired amine (6-25 mmol) and an appropriate base, like for example NaHCO_3 , K_2CO_3 , Et_3N or DIPEA (10-15 mmol) was stirred in DMSO, DMF or toluene (20-50 mL) at 60-130 °C until tlc indicated complete disappearance of the chloride. The reaction was cooled to 23 °C poured into ice-water, neutralized with 1N HCl, the precipitate was filtered off, washed with water and dried in vacuum. In cases where the product did not precipitate, the mixture was extracted with EtOAc, washed with water and brine, dried over Na_2SO_4 . Filtration and removal of the solvent in vacuum left a crude product, which was purified by silica gel column chromatography with hexane/EtOAc to give the pure title compound. Protection of the amino-group was achieved by following the general procedure B.

Example C1

[4-Iodo-5-(4-methyl-piperazin-1-yl)-2-nitro-phenyl]-carbamic acid tert.-butyl ester

Prepared from (5-chloro-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B1) (2.39 g, 6.0 mmol) and 1-methylpiperazine (1.60 mL, 15 mmol) in toluene (4.5 mL) at 110 °C for 18 h according to the general procedure C (method a). Obtained as a yellow solid (2.2 g).

MS (ISP) 463 $[(\text{M}+\text{H})^+]$; mp 134-136 °C.

Example C2

(4-Iodo-2-nitro-5-thiomorpholin-4-yl-phenyl)-carbamic acid tert.-butyl ester

Prepared from (5-chloro-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B1) (2.0 g, 5.0 mmol) and thiomorpholine (2.6 mL) in toluene (3.8 mL) DIPEA (1.7 mL) at 115 °C for 48 h according to the general procedure C (method a). Obtained as a yellow solid (1.1 g).

MS (ISP) 466 $[(\text{M}+\text{H})^+]$; mp 132-134 °C.

Example C3

(4-Iodo-5-morpholin-4-yl-2-nitro-phenyl)-carbamic acid tert.-butyl ester

Prepared from (5-chloro-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B1) (2.0 g, 5.0 mmol) and morpholine (10 mL) at reflux for 3 h according to the general procedure C (method a). Obtained as a yellow solid (0.805 g).

MS (ISP) 450 [(M+H)⁺]; mp 43-44 °C.

Example C4

[5-(1,4-Dioxo-8-aza-spiro[4.5]dec-8-yl)-4-iodo-2-nitro-phenyl]-carbamic acid tert.-butyl ester

- 5 Prepared from (5-chloro-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B1) (3.0 g, 7.53 mmol), 1,4-dioxo-8-aza-spiro(4,5)decane (4.82 mL, 37.63 mmol) and DIPEA (2.58 mL, 15.0 mmol) in toluene (4 mL) at reflux for 6 h according to the general procedure C (method a). Obtained as an orange solid (4.0 g).

MS (ISP) 506 [(M+H)⁺]; mp 132-134 °C.

10 Example C5

[4-Iodo-5-(4-methoxy-piperidin-1-yl)-2-nitro-phenyl]-carbamic acid tert.-butyl ester

- Prepared from 5-chloro-4-iodo-2-nitro-phenylamine (Example A1) (6.91 g, 23.15 mmol), 4-methoxypiperidine (4.0 g, 34.73 mmol) and NaHCO₃ (5.83 g, 69.45 mmol) in DMSO (230 mL) at 100 °C according to the general procedure C (method b). The obtained brown
15 solid (7.95 g) was converted to the title compound according to the general procedure B (method c). Obtained as a yellow solid (6.55 g).

MS (ISP) 478 [(M+H)⁺]; mp 133-135 °C.

Example C6

[5-[(2-Hydroxy-ethyl)-methyl-amino]-4-iodo-2-nitro-phenyl]-carbamic acid tert.-butyl

20 ester

Prepared from (5-chloro-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B1) (1.99 g, 5.0 mmol) and 2-methylaminoethanol (2.00 mL, 25.0 mmol) in DMSO (2.5 mL) at 23 °C according to the general procedure C (method a). Obtained as a yellow gum (1.88 g).

- 25 MS (ISP) 438 [(M+H)⁺].

Example C7

[5-(4-Hydroxy-piperidin-1-yl)-4-iodo-2-nitro-phenyl]-carbamic acid tert.-butyl ester

Prepared from (5-chloro-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B1) (1.99 g, 5 mmol) and 4-hydroxypiperidine (2.53 g, 25 mmol) in DMSO (2.5 mL) at 23 °C according to the general procedure C (method a). Obtained as a yellow solid (1.88 g).

MS (EI) 463 (M^+); mp 58-60 °C.

5 Example C8

{5-[4-(2-Hydroxy-ethoxy)-piperidin-1-yl]-4-iodo-2-nitro-phenyl}-carbamic acid tert.-butyl ester

Prepared from (5-chloro-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B1) (2.2 g, 5.5 mmol), 4-(2-hydroxyethoxy)piperidine {CAS-No. [40256-14-2]} (800 mg, 5.5 mmol) and Et_3N (2.3 mL, 16.5 mmol) in DMSO (2.3 mL) at 23 °C according to the
10 general procedure C (method a). Obtained as a yellow solid (1.65 g).

MS (EI) 507 (M^+); mp 64-65 °C.

Example C9

[5-(cis-3,4-Dihydroxy-pyrrolidin-1-yl)-4-iodo-2-nitro-phenyl]-carbamic acid tert.-butyl
15 ester

Prepared in two steps as follows: (5-chloro-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B1) (8.73 g, 21.9 mmol) was reacted with 3-pyrroline (2.0 mL, 26.3 mmol, 70% pure, contains 30% pyrrolidine), Et_3N (9.12 mL, 65.7 mmol) in DMSO (14 mL) and EtOH (5 mL) at 23 °C according to the general procedure C (method a) to give a
20 7:3 mixture of [5-(2,5-dihydro-pyrrol-1-yl)-4-iodo-2-nitro-phenyl]-carbamic acid tert.-butyl ester and (4-iodo-2-nitro-5-pyrrolidin-1-yl-phenyl)-carbamic acid tert.-butyl ester
Obtained as a yellow solid (8.57 g). Part (4.31 g) of this material was dihydroxylated by reaction with NMO (1.28 g, 11.0 mmol), 2.5% OsO_4 in t-BuOH (1 mL, 0.1 mmol) and K_2OsO_4 (40 mg, 0.1 mmol) in acetone (250 mL) and H_2O (100 mL) at 23 °C for 6 days.
25 Obtained the title compound as an amorphous yellow substance (2.50 g).

MS (ISP) 466 [$(M+H)^+$].

Example C10

[5-(2-Hydroxy-ethylamino)-4-iodo-2-nitro-phenyl]-carbamic acid tert.-butyl ester

Prepared from (5-chloro-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B1) (3.99 g, 10 mmol) and ethanolamine (3.01 mL, 50 mmol) in DMSO (20 mL) at 23 °C according to the general procedure C (method a). Obtained as a yellow solid (4.53 g).

MS (ISP) 424 [(M+H)⁺]; mp 130-148 °C.

5 Example C11

[5-((R)-3-Hydroxy-pyrrolidin-1-yl)-4-iodo-2-nitro-phenyl]-carbamic acid tert.-butyl ester

Prepared from (5-chloro-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B1), (R)-3-hydroxypyrrolidine hydrochloride and Et₃N in DMSO at 23 °C according to the general procedure C (method a). Obtained as a yellow solid (3.153 g).

10 MS (ISP) 450 [(M+H)⁺]; mp 158 °C (dec.).

The following procedures relate to the preparation of those (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl esters bearing sulfur substituents in the 5-position (Synthetic Scheme D):

General procedure D:

- 15 Preparation of 5-S-substituted-(4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl esters from (5-chloro-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester
- To a solution of the thiol (2.2 mmol) in DMF was added NaOMe-sol. (5.4M in MeOH, 0.41 mL, 2.2 mmol) followed by (5-chloro-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B1) (797 mg, 2.0 mmol) and stirring was continued at 23 °C until the
- 20 indicated complete disappearance of the chloride. Poured into ice-cold 5% citric acid, extracted with EtOAc, washed with sat. NaHCO₃-sol., brine, dried over MgSO₄. Removal of the solvent left an orange oil, which was purified by silica gel column chromatography with hexane/EtOAc to give the pure title compound.

Example D1

- 25 [5-(2-Dimethylamino-ethylsulfanyl)-4-iodo-2-nitro-phenyl]-carbamic acid tert.-butyl ester

Prepared from (5-chloro-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B1) (399 mg, 1.0 mmol), 2-dimethylaminoethanethiol hydrochloride (312 mg, 2.2 mmol) and NaOMe solution 5.4M in MeOH (0.8 mL, 8.8 mmol) in DMF (2 mL)

- 30 according to the general procedure D. Obtained as a yellow solid (306 mg).

MS (ISP) 468 [(M+H)⁺]; mp 144 °C.

Example D2

(5-tert.-Butoxycarbonylamino-2-iodo-4-nitro-phenylsulfanyl)-acetic acid methyl ester

Prepared from (5-chloro-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example
5 B1) (797 mg, 2.0 mmol), methyl thioglycolate (0.2 mL, 2.2 mmol) and NaOMe solution
5.4M in MeOH (0.41 mL, 2.2 mmol) in DMF (2 mL) according to the general procedure
D. Obtained as a yellow solid (847 mg).

MS (EI) 468 (M⁺); mp 110-112 °C.

The following procedures relate to the preparation of those (4-iodo-2-nitro-phenyl)-
10 carbamic acid tert.-butyl esters bearing oxygen substituents in the 5-position (Scheme E):

General procedure E:

Preparation of 5-O-substituted-4-iodo-2-nitro-phenylamines from 5-chloro-4-iodo-2-
nitro-phenylamine

To a suspension of KOH (85 %, 3.62-7.96 g, 55-121 mmol) in DMSO (50 mL) was added
15 the alcohol (125-500 mmol) and the mixture was stirred at 23 °C until all KOH had
dissolved. 5-Chloro-4-iodo-2-nitro-phenylamine (Example A1) (15.0 g, 50 mmol) was
added in small portions and the resulting dark red clear solution was stirred at 23-60 °C
until tlc indicated complete disappearance of the chloride. Poured into ice-cold 1N HCl or
ice-cold sat. NH₄Cl-sol., extracted with EtOAc or CHCl₃, washed with 1N HCl or sat.
20 NH₄Cl-sol. and brine, dried over MgSO₄. Removal of the solvent left a dark red solid,
which was purified by silica gel column chromatography to give the pure title compound.

Example E1

5-Allyloxy-4-iodo-2-nitro-phenylamine

Prepared from 5-chloro-4-iodo-2-nitro-phenylamine (Example A1) (15.0 g, 50 mmol),
25 allyl alcohol (50 mL) and KOH (7.96 g, 121 mmol) in DMSO (50 mL) according to the
general procedure E. Obtained as an orange solid (9.38 g).

MS (EI) 320 (M⁺); mp 74 °C.

Example E2

4-Iodo-5-methoxy-2-nitro-phenylamine

Prepared from 5-chloro-4-iodo-2-nitro-phenylamine (Example A1) (2.98 g, 10 mmol), methanol (10 mL) and KOH (1.45 g, 22 mmol) in DMSO (10 mL) according to the general procedure E. Obtained as an orange solid (2.9 g).

MS (ISP) 295 [(M+H)⁺] and 312 [(M+NH₄)⁺]; mp 189 °C.

5 Example E3

4-Iodo-5-(2-methoxy-ethoxy)-2-nitro-phenylamine

Prepared from 5-chloro-4-iodo-2-nitro-phenylamine (Example A1) (2.98 g, 10 mmol), 2-methoxyethanol (7.9 mL, 100 mmol) and KOH (1.45 g, 22 mmol) in DMSO (8 mL) according to the general procedure E. Obtained as an orange solid (2.8 g).

10 MS (ISN) 337 [(M-H)⁻]; mp 121-122 °C.

Example E4

4-Iodo-5-(2-{2-[2-(2-methoxy-ethoxy)-ethoxy]-ethoxy}-ethoxy)-2-nitro-phenylamine

Prepared from 5-chloro-4-iodo-2-nitro-phenylamine (Example A1) (9.48 g, 32 mmol), tetraethyleneglycol monomethyl ether (19 g, 91 mmol) and KOH (2.31 g, 35 mmol) in
15 DMSO (25 mL) at 60 °C according to the general procedure E. Obtained as a red oil (8.4 g).

MS (ISP) 471 [(M+H)⁺].

Example E5

(RS)-5-(2,2-Dimethyl-[1,3]dioxolan-4-ylmethoxy)-4-iodo-2-nitro-phenylamine

20 Prepared from 5-chloro-4-iodo-2-nitro-phenylamine (Example A1) (4.48 g, 15 mmol), D,L- α,β -isopropylidene-glycerol (10 mL, 81 mmol) and KOH (1.01 g, 18 mmol) in DMSO (10 mL) at 23 °C according to the general procedure E. Obtained as a yellow solid (4.9 g).

MS (ISN) 393 [(M-H)⁻]; mp 151 °C.

Example E6

25 5-(2-tert.-Butoxy-ethoxy)-4-iodo-2-nitro-phenylamine

Prepared from 5-chloro-4-iodo-2-nitro-phenylamine (Example A1) (14.9 g, 50 mmol), 2-tert.-butoxyethanol (29.5 g, 250 mmol) and KOH (3.99 g, 60 mmol) in DMSO (25 mL) at 23 °C according to the general procedure E. Obtained as a yellow solid (14.3 g).

MS (ISP) 381 [(M+H)⁺]; mp 144-146 °C.

5 Example E7

4-Iodo-5-(4-methoxy-benzyloxy)-2-nitro-phenylamine

Prepared from 5-chloro-4-iodo-2-nitro-phenylamine (Example A1) (5.97g, 20 mmol), 4-methoxybenzyl alcohol (4.98 mL, 40 mmol) and KOH (1.58 g, 24 mmol) in DMSO (30 mL) at 23 °C according to the general procedure E. Obtained as a yellow-brown solid (2.94 g).

MS (ISN) 399 [(M-H)⁻]; mp 183 °C.

The following examples relate to the preparation of (5-tert.-butoxycarbonylamino-2-iodo-4-nitro-phenyl)-acetic acid methyl ester and (5-cyanomethyl-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Synthetic Scheme F):

15 Example XVI (R=CO₂Me; R'=Me; R''=Boc)

2-(5-tert.-Butoxycarbonylamino-2-iodo-4-nitro-phenyl)-malonic acid dimethyl ester

To a solution of KOBut (0.56 g, 5.02 mmol) in DMSO (3 mL) was added dimethyl malonate (0.58 mL, 5.02 mmol) followed by (5-chloro-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B1) (1.00 g, 2.51 mmol) and the resulting dark red clear solution was stirred at 100 °C until tlc indicated complete disappearance of the chloride. Poured into ice-cold 5% citric acid (100 mL), extracted with EtOAc (2 x 100 mL), washed with brine, dried over MgSO₄. Removal of the solvent left a yellow oil, which was purified by silica gel column chromatography with hexane/EtOAc 4:1 to give the pure title compound as a yellow gum (1.13 g, 91 %).

25 MS (ISP) 512 [(M+NH₄)⁺] and 517 [(M+Na)⁺].

Example XVII (R=CN; R'=Et; R''=H)

(RS)-(5-Amino-2-iodo-4-nitro-phenyl)-cyano-acetic acid ethyl ester

Prepared as described for example XVI from 5-chloro-4-iodo-2-nitro-phenylamine (Example A1) (14.9 g, 50 mmol), ethyl cyanoacetate (14.7 mL, 100 mmol) and KOBut (11.2 g, 100 mmol) in DMSO (60 mL) at 100 °C for 2h. Obtained as a dark brown gum.

MS (EI) 375 (M^+).

5 Example Ve (R=CO₂Me; R'=Boc)

(5-tert.-Butoxycarbonylamino-2-iodo-4-nitro-phenyl)-acetic acid methyl ester

A mixture of 2-(5-tert.-butoxycarbonylamino-2-iodo-4-nitro-phenyl)-malonic acid dimethyl ester (Example XVI (R=CO₂Me; R'=Me; R''=Boc)) (3.34 g, 6.76 mmol), LiCl (573 mg, 13.52 mmol) and H₂O (0.122 mL, 6.76 mmol) in DMSO (46 mL) was stirred at
10 100 °C for 7 h. Poured into ice-water, extracted twice with EtOAc, washed with brine, dried over MgSO₄. Removal of the solvent left a yellow oil, which was purified by silica gel column chromatography with hexane/EtOAc 9:1 to give the pure title compound as a yellow solid (1.37 g, 47 %).

MS (EI) 436 (M^+); mp 93 °C.

15 Example Vf (R=CN; R''=H)

(5-Amino-2-iodo-4-nitro-phenyl)-acetonitrile

Prepared as described for example Ve from (RS)-(5-amino-2-iodo-4-nitro-phenyl)-cyanoacetic acid ethyl ester (Example XVII (R=CN; R'=Et; R''=H)) (20.62 g, 55 mmol) and LiCl (9.33 g, 220mmol) in DMSO (370 mL) and H₂O (4.4 mL) at 120 °C for 2.5 h. Obtained as a
20 green-brown solid.

MS (EI) 303 (M^+); mp 145-183 °C.

The following examples relate to the preparation of (5-hydroxymethyl-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester, the corresponding THP-ether, as well as the (5-dimethylaminomethyl-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester via the
25 intermediate chloride (Synthetic Scheme F):

Example Vh

(5-Hydroxymethyl-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester

LiBH₄ (0.32 g, 14.78 mmol) was added to a solution of 5-tert.-butoxycarbonylamino-2-iodo-4-nitro-benzoic acid methyl ester (Example B3) (3.12 g, 7.39 mmol) and MeOH (0.6 mL, 14.78 mmol) in THF (44 mL) at 0 °C. The reaction mixture was stirred at 0 °C for 15 min. Poured into 5% citric acid, extracted twice with EtOAc, washed with sat. NaHCO₃-sol. and brine, dried over MgSO₄. Removal of the solvent left a yellow oil, which was purified by silica gel column chromatography with cyclohexane/EtOAc 4:1 to give the pure title compound as a yellow solid (2.64 g, 91 %).

MS (ISP) 412 [(M+NH₄)⁺]; mp >250 °C.

Example Vi

10 (RS)-[4-Iodo-2-nitro-5-(tetrahydro-pyran-2-yloxymethyl)-phenyl]-carbamic acid tert.-butyl ester

To a mixture of (5-hydroxymethyl-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example Vh) (394 mg, 1.0 mmol) and 3,4-dihydro-2H-pyran (0.11 mL, 1.2 mmol) in DCM (5 mL) at 0 °C was added p-TsOH.H₂O (ca. 5 mg) and the reaction was stirred at 0 °C for 1 h. Diluted with EtOAc, washed with sat. NaHCO₃-sol. and brine, dried over MgSO₄. Removal of the solvent in vacuum left a yellow oil, which was purified by silica gel column chromatography with hexane/EtOAc 9:1 to give the pure title compound as a yellow gum (470 mg, 98 %).

MS (ISN) 477 [(M-H)⁻].

20 Example VI

(5-Dimethylaminomethyl-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester

To a mixture of (5-hydroxymethyl-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example Vh), LiCl (3 eq.) and pyridine (2 eq.) in DMF at 0 °C was added methanesulfonyl chloride (1.5 eq.) and the reaction was stirred at 23 °C for 24 h. Me₂NH in EtOH (10 eq.) was added and stirring was continued for 24 h. Diluted with EtOAc, washed with sat. NaHCO₃-sol. and brine, dried over MgSO₄. Removal of the solvent in vacuum left a yellow oil, which was purified by silica gel column chromatography with cyclohexane/EtOAc 3:1 to give the pure title compound as a yellow oil (421 mg).

MS (ISP) 422 [(M+H)⁺].

The following examples relate to the preparation of (2-amino-4-aryl-phenyl)-carbamic acid tert.-butyl esters, [2-amino-4-(1-alkenyl)-phenyl]-carbamic acid tert.-butyl esters and (2-amino-4-aryl-phenyl)-carbamic acid tert.-butyl esters in regioisomerically pure fashion (Synthetic Scheme B):

5 General procedure F:

- Preparation of (4-aryl-2-nitro-phenyl)-carbamic acid tert.-butyl esters by direct Suzuki-coupling of (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl esters with arylboronic acids. A mixture of the (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (3.0 mmol), the arylboronic acid (4.5 mmol) and $\text{PdCl}_2(\text{PPh}_3)_2$ (2 mol%) was refluxed in 1,4-dioxane (25 mL) and 2M Na_2CO_3 -sol. (7.5 mL) [or alternatively with 1M NaHCO_3 -sol. (7.5 mL), LiCl (6.0 mmol) and $(\text{Ph}_3\text{P})_4\text{Pd}$ (3 mol%) in DME (30 mL); also possible with Et_3N (9.0 mmol), $\text{Pd}(\text{OAc})_2$ (3 mol%), PPh_3 (6 mol%) in DMF (10 mL) at 100 °C] until tlc indicated complete conversion of the iodide. The mixture was transferred into a separating funnel, H_2O (25 mL) was added and the product was extracted with ether or EtOAc (3 x 30 mL). The combined organic layers were washed with brine (50 mL) and dried over Na_2SO_4 . Removal of the solvent left a brown residue, which was purified by silica gel column chromatography with cyclohexane/ether or cyclohexane/EtOAc to give the title compound.

Example F1

(2-Chloro-5-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester

- 20 Prepared from (5-chloro-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B1) (1.20 g, 3.00 mmol) and phenyl boronic acid (0.62 g, 3.30 mmol) according to the general procedure F. Obtained as a yellow oil (843 mg).

MS (EI) 348 (M^+) and 350 [$(\text{M}+2)^+$].

Example F2

- 25 (2-Methyl-5-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (4-iodo-5-methyl-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B2) (1.135 g, 3 mmol) and phenylboronic acid (630 mg, 3.3 mmol) according to the general procedure F. Obtained as a yellow oil (971 mg).

MS (EI) 328 (M^+).

Example F3

(RS)-{4'-Fluoro-5-nitro-2-[4-(tetrahydro-pyran-2-yloxy)-piperidin-1-yl]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (RS)-{4-iodo-2-nitro-5-[4-(tetrahydro-pyran-2-yloxy)-piperidin-1-yl]-phenyl}-carbamic acid tert.-butyl ester [RO-69-4319/000, prepared from [5-(4-hydroxy-piperidin-1-yl)-4-iodo-2-nitro-phenyl]-carbamic acid tert.-butyl ester (Example C7) by treatment with 3,4-dihydro-2H-pyran and cat. TsOH.H₂O in DCM at 0 °C] (1.09 g, 2.0 mmol) and 4-fluorophenylboronic acid according to the general procedure F. Obtained as an orange solid (894 mg).

MS (ISP) 516 [(M+H)⁺]; mp 144-146 °C.

Example F4

(RS)-(4'-Fluoro-5-nitro-2-[4-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-piperidin-1-yl]-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (RS)-(4-iodo-2-nitro-5-[4-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-piperidin-1-yl]-phenyl)-carbamic acid tert.-butyl ester [RO-69-4355/000, prepared from {5-[4-(2-hydroxy-ethoxy)-piperidin-1-yl]-4-iodo-2-nitro-phenyl}-carbamic acid tert.-butyl ester (Example C8) by treatment with 3,4-dihydro-2H-pyran and cat. TsOH.H₂O in DCM at 0 °C] (950 mg, 1.87 mmol) and 4-fluorophenylboronic acid (314 mg, 2.25 mmol) according to the general procedure F. Obtained as a viscous orange oil (930 mg).

MS (ISP) 560 [(M+H)⁺]; mp 144-146 °C.

Example F5

(RS)-[4'-Fluoro-5-nitro-2-(tetrahydro-pyran-2-yloxymethyl)-biphenyl-4-yl]-carbamic acid tert.-butyl ester

Prepared from (RS)-[4-iodo-2-nitro-5-(tetrahydro-pyran-2-yloxymethyl)-phenyl]-carbamic acid tert.-butyl ester (Example Vi) and 4-fluorophenylboronic acid according to the general procedure F. Obtained as an orange oil (1.24 g).

Example F6

(2-Cyanomethoxy-4'-fluoro-5-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (5-cyanomethoxy-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example S2) (838 mg, 2.0 mmol) and 4-fluorophenylboronic acid (392 mg, 2.8 mmol) according to the general procedure F. Obtained as a yellow solid (333 mg).

MS (ISP) 405 [(M+NH₄)⁺]; mp 148 °C.

5 Example F7

(2-Dimethylaminomethyl-4'-fluoro-5-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (5-dimethylaminomethyl-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example VI) and 4-fluorophenylboronic acid according to the general procedure F. Obtained as a yellow solid (1.01 g).

10 Example F8

[2-(2,2-Dimethyl-tetrahydro-[1,3]dioxolo[4,5-c]pyrrol-5-yl)-4'-fluoro-5-nitro-biphenyl-4-yl]-carbamic acid tert.-butyl ester

Prepared from [5-(cis-2,2-dimethyl-tetrahydro-[1,3]dioxolo[4,5-c]pyrrol-5-yl)-4-iodo-2-nitro-phenyl]-carbamic acid tert.-butyl ester [RO-69-4741/000, prepared from [5-(cis-3,4-dihydroxy-pyrrolidin-1-yl)-4-iodo-2-nitro-phenyl]-carbamic acid tert.-butyl ester
15 (Example C9) by treatment with 2,2-dimethoxypropane and cat. TsOH.H₂O in DMF at 23 °C] (845 mg, 1.67 mmol) and 4-fluorophenylboronic acid (327 mg, 2.34 mmol) according to the general procedure F. Obtained as a yellow solid (643 mg).

MS (ISP) 474 [(M+H)⁺]; mp 119 °C.

20 Example F9

(4'-Fluoro-2-methoxy-5-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (4-iodo-5-methoxy-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B5) (3.68 g, 9.34 mmol) and 4-fluorophenylboronic acid (3.61 g, 25.8 mmol) according to the general procedure F. Obtained as a yellow solid (2.69 g).

25 MS (ISN) 361 [(M-H)⁻]; mp 250 °C.

Example F10

[2-(1,4-Dioxo-8-aza-spiro[4.5]dec-8-yl)-4'-fluoro-5-nitro-biphenyl-4-yl]-carbamic acid tert.-butyl ester

Prepared from [5-(1,4-dioxo-8-aza-spiro[4.5]dec-8-yl)-4-iodo-2-nitro-phenyl]-carbamic acid tert.-butyl ester (Example C4) (4.0 g, 7.02 mmol) and 4-fluorophenylboronic acid (1.33 g, 9.5 mmol) according to the general procedure F. Obtained as a yellow solid (2.43 g).

5 mp 213 °C (dec.).

Example F11

(4'-Fluoro-2-methyl-5-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (4-iodo-5-methyl-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B2) (756 mg, 2.0 mmol) and 4-fluorophenylboronic acid (420 mg, 3.0 mmol) according to
10 the general procedure F. Obtained as an amorphous yellow substance (611 mg).

MS (ISN) 345 [(M-H)].

Example F12

(4-tert.-Butoxycarbonylamino-4'-fluoro-5-nitro-biphenyl-2-yloxy)-acetic acid tert.-butyl ester

15 Prepared from (5-tert.-butoxycarbonylamino-2-iodo-4-nitro-phenoxy)-acetic acid tert.-butyl ester (Example S1) (2.14 g, 4.33 mmol) and 4-fluorophenylboronic acid (728 mg, 5.2 mmol) according to the general procedure F. Obtained as an orange solid (1.80 g).

MS (ISN) 461 [(M-H)]; mp 92-93 °C.

Example F13

20 (2-Chloro-4'-fluoro-5-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (5-chloro-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B1) and 4-fluorophenylboronic acid according to the general procedure F. Obtained as a yellow solid (625 mg).

MS (EI) 366 (M⁺).

25 Example F14

[4'-Fluoro-2-(2-methoxy-ethoxy)-5-nitro-biphenyl-4-yl]-carbamic acid tert.-butyl ester

Prepared from [4-iodo-5-(2-methoxy-ethoxy)-2-nitro-phenyl]-carbamic acid tert.-butyl ester (Example B6) and 4-fluorophenylboronic acid according to the general procedure F. Obtained as a yellow solid (1.833 g).

MS (EI) 406 (M^+).

5 Example F15

[2-(2-tert.-Butoxy-ethoxy)-4'-fluoro-5-nitro-biphenyl-4-yl]-carbamic acid tert.-butyl ester

Prepared from [5-(2-tert.-butoxy-ethoxy)-4-iodo-2-nitro-phenyl]-carbamic acid tert.-butyl ester (Example B10) and 4-fluorophenylboronic acid according to the general procedure F. Obtained as a yellow solid (735 mg).

10 MS (ISP) 449 [$(M+H)^+$].

Example F16

[4'-Fluoro-5-nitro-2-(2-oxo-oxazolidin-3-yl)-biphenyl-4-yl]-carbamic acid tert.-butyl ester

Prepared from [4-iodo-2-nitro-5-(2-oxo-oxazolidin-3-yl)-phenyl]-carbamic acid tert.-butyl ester [RO-69-6758/000, prepared from [5-(2-hydroxy-ethylamino)-4-iodo-2-nitro-phenyl]-carbamic acid tert.-butyl ester (Example C10) by treatment with 1,1'-carbonyldiimidazole in dioxane and then in pyridine with cat. DMAP each at 100 °C] (503 mg, 1.12 mmol) and 4-fluorophenylboronic acid (235 mg, 1.68 mmol) according to the general procedure F. Obtained as a yellow solid (310 mg).

20 MS (ISN) 416 [$(M-H)^+$]; mp 201 °C.

Example F17

(4'-Fluoro-2-methoxy-2'-methyl-5-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (4-iodo-5-methoxy-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B5) and 4-fluoro-2-methyl-phenylboronic acid according to the general procedure F. Obtained as a yellow solid (699 mg).

25 MS (EI) 376 (M^+).

Example F18

(2-tert.-Butoxy-4'-fluoro-5-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (5-tert.-butoxy-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester

(Example S4) (1.4 g, 3.21 mmol) and 4-fluorophenylboronic acid (0.67 g, 4.42 mmol)

5 according to the general procedure F. Obtained as an amorphous yellow substance (1.2 g).

MS (EI) 404 (M^+).

Example F19

(2-tert.-Butoxy-2'-fluoro-5-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (5-tert.-butoxy-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester

10 (Example S4) (1.4 g, 3.21 mmol) and 2-fluorophenylboronic acid (0.67 g, 4.83 mmol)

according to the general procedure F. Obtained as an amorphous yellow substance (960 mg).

MS (EI) 404 (M^+).

Example F20

15 (RS)-{4'-Fluoro-5-nitro-2-[(R)-3-(tetrahydro-pyran-2-yloxy)-pyrrolidin-1-yl]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (RS)-{4-iodo-2-nitro-5-[(R)-3-(tetrahydro-pyran-2-yloxy)-pyrrolidin-1-

yl]-phenyl}-carbamic acid tert-butyl ester [RO-69-6376/000, prepared from [5-((R)-3-

20 hydroxy-pyrrolidin-1-yl)-4-iodo-2-nitro-phenyl]-carbamic acid tert.-butyl ester (Example C11) by treatment with 3,4-dihydro-2H-pyran and cat. TsOH.H₂O in DCM at 0 °C] and 4-fluorophenylboronic acid according to the general procedure F. Obtained as a yellow solid (1.053 g).

MS (ISP) 502 [(M+H)⁺].

Example F21

25 (2'-Fluoro-2-methoxy-5-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (4-iodo-5-methoxy-2-nitro-phenyl)-carbamic acid tert.-butyl ester

(Example B5) (1.00 g, 2.54 mmol) and 2-fluorophenylboronic acid (0.60 g, 4.32 mmol)

according to the general procedure F. Obtained as an amorphous yellow substance (687 mg).

MS (EI) 362 (M^+).

Example F22

- 5 [2-(1,4-Dioxo-8-aza-spiro[4.5]dec-8-yl)-2'-fluoro-5-nitro-biphenyl-4-yl]-carbamic acid tert.-butyl ester

Prepared from [5-(1,4-dioxo-8-aza-spiro[4.5]dec-8-yl)-4-iodo-2-nitro-phenyl]-carbamic acid tert.-butyl ester (Example C4) (1.89 g, 3.47 mmol) and 2-fluorophenylboronic acid (0.63 g, 4.49 mmol) according to the general procedure F. Obtained as a yellow solid (1.46 g).

MS (ISP) 474 [($M+H$) $^+$]; mp 164 °C.

Example F23

(2',5'-Difluoro-2-methoxy-5-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester

- Prepared from (4-iodo-5-methoxy-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B5) (3.94 g, 10 mmol) and 2,5-difluorophenylboronic acid (2.21 g, 14 mmol) according to the general procedure F. Obtained as an amorphous yellow substance (1.05 g).

MS (ISN) 379 [($M-H$) $^-$].

Example F24

- 20 [2'-Fluoro-2-(2-methoxy-ethoxy)-5-nitro-biphenyl-4-yl]-carbamic acid tert.-butyl ester

Prepared from [4-iodo-5-(2-methoxy-ethoxy)-2-nitro-phenyl]-carbamic acid tert.-butyl ester (Example B6) and 2-fluorophenylboronic acid according to the general procedure F. Obtained as a yellow solid (3.63 g).

MS (ISN) 405 [($M-H$) $^-$].

- 25 Example F25

(RS)-[2'-Fluoro-5-nitro-2-(tetrahydro-pyran-2-ylloxymethyl)-biphenyl-4-yl]-carbamic acid tert.-butyl ester

Prepared from (RS)-[4-iodo-2-nitro-5-(tetrahydro-pyran-2-yloxymethyl)-phenyl]-carbamic acid tert.-butyl ester (Example Vi) and 2-fluorophenylboronic acid according to the general procedure F. Obtained as a yellow liquid (2.606 g).

MS (ISN) 445 [(M-H)⁺].

5 Example F26

[2'-Fluoro-2-(4-methoxy-benzyloxy)-5-nitro-biphenyl-4-yl]-carbamic acid tert.-butyl ester

Prepared from [4-iodo-5-(4-methoxy-benzyloxy)-2-nitro-phenyl]-carbamic acid tert.-butyl ester (Example B11) (1.69 g, 3.38 mmol) and 2-fluorophenylboronic acid (0.61 g,
10 4.39 mmol) according to the general procedure F. Obtained as a yellow foam (940 mg).

MS (ISP) 469 [(M+H)⁺].

Example F27

(2-tert.-Butoxy-2',5'-difluoro-5-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (5-tert.-butoxy-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester
15 (Example S4) (3.00 g, 6.88 mmol) and 2,5-difluorophenylboronic acid (2.23 g, 14.1 mmol) according to the general procedure F. Obtained as an amorphous yellow substance (2.30 g).

MS (ISN) 421 [(M-H)⁺].

General procedure G:

20 Preparation of (4-aryl-2-nitro-phenyl)-carbamic acid tert.-butyl esters by Suzuki-coupling of (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl esters with bis(pinacolato)diboron and subsequent reaction with aryl halides

A mixture of the (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (2.0 mmol), bis(pinacolato)diboron (2.2 mmol), KOAc (6.0 mmol) and PdCl₂(PPh₃)₂ (3 mol%) in 1,4-dioxane (25 mL) was stirred at 100 °C until tlc indicated complete conversion of the iodide
25 [cf. *Tetr. Lett.* 1997, 38, 3841-3844]. After addition of the aryl halide (4.0 mmol), PdCl₂(PPh₃)₂ (3 mol%) and 2M Na₂CO₃-sol. (7.5 mL) the mixture was stirred at 100 °C until tlc indicated complete conversion of the intermediate boronic ester. The mixture was transferred into a separating funnel, H₂O (30 mL) was added and the product was
30 extracted with ether or EtOAc (3 x 50 mL). The combined organic layers were washed with

brine (100 mL) and dried over Na_2SO_4 . Removal of the solvent left a brown residue, which was purified by silica gel column chromatography with cyclohexane/ether or cyclohexane/EtOAc to give the title compound.

General procedure H:

- 5 Preparation of (4-aryl-2-nitro-phenyl)-carbamic acid tert.-butyl esters by carbonylative Suzuki-coupling of (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl esters with aryl boronic acids
- A mixture of the (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (1.0 mmol), aryl boronic acid (1.1 mmol), K_2CO_3 (3.0 mmol) and $\text{PdCl}_2(\text{PPh}_3)_2$ (3 mol%) in anisole (6 mL)
- 10 was stirred at 80 °C under a CO-atmosphere until thin layer chromatography indicated complete conversion of the iodide [cf. *Tetr. Lett.* 1993, 34, 7595-7598]. The mixture was transferred into a separating funnel, H_2O (30 mL) was added and the product was extracted with EtOAc (2×100 mL). The combined organic layers were washed with brine (50 mL) and dried over Na_2SO_4 . Removal of the solvent left a yellow residue, which was
- 15 purified by silica gel column chromatography with or hexane/EtOAc to give the title compound.

General procedure I:

- Preparation of (4-aryl-2-nitro-phenyl)-carbamic acid tert.-butyl esters or (4-{alkenyl-, cycloalkenyl- or heterocycloalkenyl}-2-nitro-phenyl)-carbamic acid tert.-butyl esters by
- 20 Stille-coupling of (2-nitro-4-tributylstannanyl-phenyl)-carbamic acid tert.-butyl ester with aryl halides or vinyl triflates or Stille-coupling of (4-iodo-2-nitrophenyl)-carbamic acid tert.-butyl ester with trialkylaryl stannanes
- A mixture of (2-nitro-4-tributylstannanyl-phenyl)-carbamic acid tert.-butyl ester (525 mg, 1.0 mmol; prepared from the corresponding (4-iodo-2-nitro-phenyl)-carbamic acid tert.-
- 25 butyl ester (Examples B) (10 mmol) by reaction with hexabutyldistannane (7.5 mL, 15 mmol) and $\text{Pd}(\text{PPh}_3)_4$ (116 mg, 0.1 mmol) in toluene (20 mL) at 60 °C for 5 days according to *Bull. Chem. Soc. Jpn.* 1983, 56, 3855-3856), aryl halide or vinyl triflate (0.95-6.0 mmol), anhydrous LiCl (126 mg, 3.0 mmol) and $\text{Pd}(\text{PPh}_3)_4$ (5 mol%) in DME (3 mL) was stirred at 100 °C under argon atmosphere until tlc indicated complete consumption of
- 30 the stannane. The reaction was cooled to 23 °C, stirred with sat. aqueous KF-sol. (5 mL) for 45 min, filtered through celite, washed with ether and the filtrate was dried over MgSO_4 . Removal of the solvent in vacuum left a brown oil, which was purified by silica gel column chromatography with hexane/EtOAc to give the title compound.

The following examples relate to the preparation of (2-amino-4-arylethynyl-phenyl)-carbamic acid tert.-butyl esters in regioisomerically pure fashion (Synthetic Scheme G):

General procedure K:

- Preparation of (4-alkynyl-2-nitro-phenyl)-carbamic acid tert.-butyl esters by Sonogashira-coupling of (4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl esters with acetylenic compounds;
also Sonogashira-coupling of (4-ethynyl-2-nitro-phenyl)-carbamic acid tert.-butyl esters with aryl halides;
and Sonogashira-coupling of 8-iodo-4-aryl-1,3-dihydro-benzo[b][1,4]diazepin-2-ones with acetylenic compounds.
- A mixture of the halide (3.0-4.5 mmol), acetylenic compound (3.0-4.5 mmol), Et₃N (13.5 mmol), PdCl₂(PPh₃)₂ (5 mol%) and PPh₃ (2.5 mol%) in THF (12 mL) [with very insoluble material DMF (up to 12 mL) could be added] was stirred for 20 min at 23 °C while being purged with Argon. CuI (1.2 mol%) was added and stirring was continued at 60 °C under Argon atmosphere until tlc indicated complete conversion of the minor component [cf. *J. Org. Chem.* 1998, 63, 8551]. The mixture was transferred into a separating funnel, 5% citric acid (50 mL) was added and the product was extracted with EtOAc (2 × 100 mL). The combined organic layers were washed with sat. NaHCO₃-sol. (50 mL) and brine (50 mL), followed by drying over MgSO₄. Removal of the solvent left a yellow residue, which was purified by silica gel column chromatography with hexane/EtOAc and/or triturated with hexane or aqueous EtOH to give the title compound.

Example K1

(5-Chloro-2-nitro-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester

- Prepared from (5-chloro-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B1) (1.2 g, 3.0 mmol) and phenylacetylene (0.5 mL, 4.5 mmol) according to the general procedure K. Obtained as a yellow solid (944 mg).

MS (ISN) 371 [(M-H)⁻] and 373 [(M-H+2)⁻]; mp 166-167 °C.

Example K2

(5-Methyl-2-nitro-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester

Prepared from (4-iodo-5-methyl-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B2) (1.13 g, 3.0 mmol) and phenylacetylene (0.5 mL, 4.5 mmol) according to the general procedure K. Obtained as a green-yellow solid (794 mg).

MS (EI) 352 (M^+); mp 161-164 °C.

5 Example K3

[5-(4-Methyl-piperazin-1-yl)-2-nitro-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester

Prepared from [4-iodo-5-(4-methyl-piperazin-1-yl)-2-nitro-phenyl]-carbamic acid tert.-butyl ester (Example C1) (1.34 g, 3.0 mmol) and phenylacetylene (0.5 mL, 4.5 mmol) according to the general procedure K. Obtained as a green-yellow solid (1.1 g).

MS (ISP) 437 [$(M+H)^+$]; mp 170 °C.

Example K4

(5-Morpholin-4-yl-2-nitro-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester

Prepared from (4-iodo-5-morpholin-4-yl-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example C3) (890 mg, 2.0 mmol) and phenylacetylene (0.33 mL, 3.0 mmol) according to the general procedure K. Obtained as an orange solid (580 mg).

MS (ISP) 424 [$(M+H)^+$]; mp 190-191 °C.

Example K5

(2-Nitro-4-phenylethynyl-5-thiomorpholin-4-yl-phenyl)-carbamic acid tert.-butyl ester

Prepared from (4-iodo-2-nitro-5-thiomorpholin-4-yl-phenyl)-carbamic acid tert.-butyl ester (Example C2) (1.0 g, 2.15 mmol) and phenylacetylene (0.36 mL, 3.22 mmol) according to the general procedure K. Obtained as an orange solid (620 mg).

MS (ISP) 440 [$(M+H)^+$] and 462 [$(M+Na)^+$]; mp 201 °C (dec.).

Example K6

[5-(1,1-Dioxo-thiomorpholin-4-yl)-2-nitro-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester

Prepared in two steps as followed:

- To a solution of (4-iodo-2-nitro-5-thiomorpholin-4-yl-phenyl)-carbamic acid tert.-butyl ester (Example C2) (465 mg, 1 mmol) in acetone (25 mL) and H₂O (1 mL) 0.3M ammoniummolybdate sol. (0.3 mL) and 33% H₂O₂ (2.3 mL) were added at 0°C and mixture was stirred for 1 h at 23 °C. Obtained the [5-(1,1-dioxo-thiomorpholin-4-yl)-4-iodo-2-nitro-phenyl]-carbamic acid tert.-butyl ester (497 mg, 1.0 mmol) as an amorphous yellow material, which was reacted with phenylacetylene (0.17 mL, 1.5 mmol) according to the general procedure K. Obtained as a yellow solid (245 mg).

MS (ISP) 472 [(M+H)⁺]; mp 217-221 °C.

10 Example K7

[5-(1,4-Dioxa-8-aza-spiro[4.5]dec-8-yl)-2-nitro-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester

- Prepared from [5-(1,4-dioxa-8-aza-spiro[4.5]dec-8-yl)-4-iodo-2-nitro-phenyl]-carbamic acid tert.-butyl ester (Example C4) (3.80 g, 7.53 mmol) and phenylacetylene (1.24 mL, 11.3 mmol) according to the general procedure K. Obtained as a orange solid (1.8 g).

MS (ISN) 478 [(M-H)⁻]; mp 179-180 °C.

Example K8

[5-(2-Dimethylamino-ethylsulfanyl)-2-nitro-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester

- 20 Prepared from [5-(2-dimethylamino-ethylsulfanyl)-4-iodo-2-nitro-phenyl]-carbamic acid tert.-butyl ester (Example D1) (721 mg, 1.54 mmol) and phenylacetylene (0.25 mL, 2.31 mmol) according to the general procedure K. Obtained as an amorphous yellow material (595 mg).

MS (ISP) 442 [(M+H)⁺]; mp 179-180 °C.

25 Example K9

(5-tert.-Butoxycarbonylamino-4-nitro-2-phenylethynyl-phenylsulfanyl)-acetic acid methyl ester

Prepared from (5-tert.-butoxycarbonylamino-2-iodo-4-nitro-phenylsulfanyl)-acetic acid methyl ester (Example D2) (780 mg, 1.67 mmol) and phenylacetylene (0.27 mL, 2.5 mmol) according to the general procedure K. Obtained as an orange solid (700 mg).

MS (ISP) 460 [(M+NH₄)⁺]; mp 125-127 °C.

5 Example K10

[5-(2-Methoxy-ethoxy)-2-nitro-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester

Prepared from [4-iodo-5-(2-methoxy-ethoxy)-2-nitro-phenyl]-carbamic acid tert.-butyl ester (Example B6) (876 mg, 2 mmol) and phenylacetylene (0.33 mL, 3 mmol) according to the general procedure K. Obtained as a yellow solid (569 mg).

10 MS (ISP) 413 [(M+H)⁺] and 430 [(M+NH₄)⁺]; mp 118-119 °C.

Example K11

(5-Methoxy-2-nitro-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester

Prepared from (4-iodo-5-methoxy-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B5) (1.18 g, 3.00 mmol) and phenylacetylene (0.58 mL, 4.5 mmol) according to
15 the general procedure K. Obtained as a yellow solid (1.1 g).

MS (EI) 368 (M⁺); mp 129 °C.

Example K12

[5-(2-{2-[2-(2-Methoxy-ethoxy)-ethoxy]-ethoxy}-ethoxy)-2-nitro-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester

20 Prepared from [4-iodo-5-(2-[2-[2-(2-methoxy-ethoxy)-ethoxy]-ethoxy]-ethoxy)-2-nitro-phenyl]-carbamic acid tert.-butyl ester (Example B7) (5.7 g, 10.0 mmol) and phenylacetylene (1.65 mL, 15 mmol) according to the general procedure K. Obtained as a yellow oil (5.2 g).

MS (ISN) 543 [(M-H)⁻].

25 Example K13

(5-tert.-Butoxycarbonylamino-4-nitro-2-phenylethynyl-phenoxy)-acetic acid tert.-butyl ester

Prepared from (5-tert.-butoxycarbonylamino-2-iodo-4-nitro-phenoxy)-acetic acid tert.-butyl ester (Example S1) (1.46 g, 2.99 mmol) and phenylacetylene (0.49 mL, 4.49 mmol) according to the general procedure K. Obtained as a yellow solid (1.4 g).

MS (ISP) 486 [(M+NH₄)⁺]; mp 130 °C.

5 Example K14

(RS)-5-(2,2-Dimethyl-[1,3]dioxolan-4-ylmethoxy)-2-nitro-4-phenylethynyl-phenylamine

Prepared from (RS)-5-(2,2-dimethyl-[1,3]dioxolan-4-ylmethoxy)-4-iodo-2-nitro-phenylamine (Example E5) (4.5 g, 11.4 mmol) and phenylacetylene (1.88 mL, 17.1 mmol) according to the general procedure K. Obtained as an orange solid (5.4 g).

10 MS (ISN) 367 [(M-H)⁻]; mp 147-149 °C.

Example K15

(5-tert.-Butoxycarbonylamino-4-nitro-2-phenylethynyl-benzoic acid methyl ester

Prepared from 5-tert.-butoxycarbonylamino-2-iodo-4-nitro-benzoic acid methyl ester (Example B3) (1.22 g, 2.89 mmol) and phenylacetylene (0.48 mL, 4.34 mmol) according to
15 the general procedure K. Obtained as a yellow solid (793 mg).

MS (ISP) 397 [(M+H)⁺] and 414 [(M+NH₄)⁺]; mp 173 °C.

Example K16

(5-tert.-Butoxycarbonylamino-4-nitro-2-phenylethynyl-phenyl)-acetic acid methyl ester

Prepared from (5-tert.-butoxycarbonylamino-2-iodo-4-nitro-phenyl)-acetic acid methyl
20 ester (Example Ve (R=CO₂Me; R''=Boc)) (1.32 g, 3.03 mmol) and phenylacetylene (0.5 mL, 4.55 mmol) according to the general procedure K. Obtained as a yellow solid (1.1 g).

MS (ISP) 411 [(M+H)⁺], 428 [(M+NH₄)⁺] and 433 [(M+Na)⁺]; mp 134 °C.

Example K17

(5-Cyanomethyl-2-nitro-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester

Prepared from (5-cyanomethyl-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B9 [Ve (R=CN; Rⁿ=Boc)]) (3.97 g, 9.85 mmol) and phenylacetylene (3.24 mL, 29.56 mmol) according to the general procedure K. Obtained as an olive solid (1.6 g).

MS (ISP) 395 [(M+NH₄)⁺]; mp 166 °C.

5 Example K18

(RS)-[5-(2,3-Dihydroxy-propoxy)-2-nitro-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester.

Prepared from (RS)-[5-(2,3-dihydroxy-propoxy)-4-iodo-2-nitro-phenyl]-carbamic acid tert.-butyl ester (RO-68-5451/000 (3.23 g, 7.11 mmol); prepared from (5-allyloxy-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B4) (4.20 g, 10.0 mmol) by reaction with NMO (1.28 g, 11.0 mmol), OsO₄ 2.5% in t-BuOH (1 mL, 0.1 mmol) and K₂OsO₄ (40 mg, 0.1 mmol) in acetone (250 mL) and H₂O (100 mL) at 23 °C for 6 days) and phenylacetylene (1.17 mL, 10.67 mmol) according to the general procedure K. Obtained as a yellow solid (2.74 g).

15 MS (ISP) 429 [(M+H)⁺]; mp 157 °C (dec.).

Example K19

(5-Hydroxymethyl-2-nitro-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (5-hydroxymethyl-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example Vh) (2.61 g, 6.62 mmol) and phenylacetylene (1.10 mL, 9.93 mmol) according to the general procedure K. Obtained as a yellow solid (1.76 g).

MS (ISP) 386 [(M+NH₄)⁺]; mp 177 °C (dec.).

Example K20

[5-(4-Methoxy-piperidin-1-yl)-2-nitro-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester.

25 Prepared from [4-iodo-5-(4-methoxy-piperidin-1-yl)-2-nitro-phenyl]-carbamic acid tert.-butyl ester (Example C5) (1.0 g, 2.1 mmol) and phenylacetylene (0.35 mL, 3.15 mmol) according to the general procedure K. Obtained as a yellow solid (799 mg).

mp 147-150 °C.

Example K21

(5-Cyanomethoxy-2-nitro-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (5-cyanomethoxy-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example S2) (605 mg, 1.44 mmol) and phenylacetylene (0.24 mL, 2.16 mmol) according to the general procedure K. Obtained as a yellow solid (508 mg).

MS (EI) 393 (M^+); mp 170 °C.

Example K22

[4-(4-Fluoro-phenylethynyl)-5-hydroxymethyl-2-nitro-phenyl]-carbamic acid tert.-butyl ester.

10 Prepared from (5-hydroxymethyl-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example Vh) (2.00 g, 5.07 mmol) and 4-fluorophenylacetylene (0.91 g, 7.61 mmol) according to the general procedure K. Obtained as a yellow solid (1.55 g).

MS (ISN) 385 [(M-H)⁻]; mp 198 °C.

Example K23

15 (RS)-(4-(4-Fluoro-phenylethynyl)-5-{methyl-[2-(tetrahydro-pyran-2-yloxy)-ethyl]-amino}-2-nitro-phenyl)-carbamic acid tert.-butyl ester.

Prepared from (RS)-(4-iodo-5-{methyl-[2-(tetrahydro-pyran-2-yloxy)-ethyl]-amino}-2-nitro-phenyl)-carbamic acid tert.-butyl ester [RO-69-3820/000, prepared from {5-[(2-hydroxy-ethyl)-methyl-amino]-4-iodo-2-nitro-phenyl}-carbamic acid tert.-butyl ester (Example C6) by treatment with 3,4-dihydro-2H-pyran and cat. TsOH.H₂O in DCM at 0 °C] (2.09 g, 4.01 mmol) and 4-fluorophenylacetylene (0.72 g, 6.02 mmol) according to the general procedure K. Obtained as a yellow-brown solid (1.84 g).

MS (ISP) 514 [(M+H)⁺]; mp 134 °C.

Example K24

25 (RS)-{2-Nitro-4-phenylethynyl-5-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-phenyl}-carbamic acid tert.-butyl ester.

Prepared from (RS)-{4-iodo-2-nitro-5-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-phenyl}-carbamic acid tert.-butyl ester (Example S3) (743 mg, 1.46 mmol) and phenylacetylene

(0.24 mL, 2.19 mmol) according to the general procedure K. Obtained as a yellow-brown viscous oil (429 mg).

MS (EI) 393 (M^+).

Example K25

- 5 Prepared from [5-(2-tert.-butoxy-ethoxy)-4-iodo-2-nitro-phenyl]-carbamic acid tert.-butyl ester (Example B10) (1.44 g, 3.0 mmol) and 4-fluorophenylacetylene (541 mg, 4.5 mmol) according to the general procedure K. Obtained as a yellow solid (777 mg).

MS (EI) 472 (M^+); mp 96-98 °C.

Example K26

- 10 [5-tert.-Butoxy-4-(4-fluoro-phenylethynyl)-2-nitro-phenyl]-carbamic acid tert.-butyl ester.

Prepared from (5-tert.-butoxy-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example S4) (1.40 g, 3.21 mmol) and 4-fluorophenylacetylene (0.66 g, 5.46 mmol) according to the general procedure K. Obtained as a brown solid (520 mg).

- 15 MS (EI) 428 (M^+); mp 201 °C.

General procedure L:

Preparation of the (2-amino-phenyl)-carbamic acid tert.-butyl esters by reduction of (2-nitro-phenyl)-carbamic acid tert.-butyl esters;

- Also preparation of 4-aryl-1,3-dihydro-benzo[b][1,4]diazepin-2-ones by reduction and
20 concomitant cyclization of 3-aryl-N-(2-nitro-phenyl)-3-oxo-propionamides

Method a: Catalytic hydrogenation

- A mixture of the nitro compound (1.0 mmol) in MeOH or EtOH and THF (1:1 ca. 20 mL) and 10% Palladium on carbon (20 mg) or Raney-Ni (20 mg) was stirred vigorously at 23
25 °C under hydrogen atmosphere until tlc indicated complete conversion. The catalyst was filtered off, washed thoroughly with MeOH or EtOH and THF (1:1), the solvent was removed in vacuum to give the title compound, which was generally pure enough for further transformations.

Method b: Reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$

- A mixture of the nitro compound (1.0 mmol) and $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (5.0 mmol) was either stirred in EtOH (30 mL) at 70-80 °C or alternatively in pyridine (3 mL) and DMF (12 mL) at 23 °C under Argon atmosphere until tlc indicated complete conversion [cf. *Tetr. Lett.* 1984, 25, 839]. The reaction mixture was brought to pH 8 by addition of sat. NaHCO_3 -sol. and extracted with EtOAc (2 x 100 mL). The combined organic layer were washed with brine and dried over Na_2SO_4 . Removal of the solvent left a yellow solid, which - if necessary - can be purified by silica gel column chromatography.

Method c: Reduction with Zn and NH_4Cl

- To a mixture of the nitro compound (1.0 mmol) in EtOH/THF/sat. NH_4Cl -sol. (1:1:1, 30 mL) was added Zinc dust (3.0 mmol) and the mixture was stirred at 70 °C under Argon atmosphere until tlc indicated complete conversion. Aqueous workup as described in method b.

Method d: Reduction with Fe and HOAc

- To a mixture of the nitro compound (1.0 mmol) in THF/ H_2O (4:1, 10-50 mL) was added Fe powder (6.0 mmol) and the mixture was stirred at 70 °C under Argon atmosphere until tlc indicated complete conversion. Aqueous workup as described in method b.

Example L1

(2-Amino-4-iodo-5-thiomorpholin-4-yl-phenyl)-carbamic acid tert.-butyl ester

- Prepared from (4-iodo-2-nitro-5-thiomorpholin-4-yl-phenyl)-carbamic acid tert.-butyl ester (Example C2) (1.05 g, 2.25 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (2.54 g, 11.3 mmol) according to the general procedure L (method b). Obtained as a light yellow solid (993 mg).

MS (ISP) 436 $[(\text{M}+\text{H})^+]$; mp 125-127 °C.

25 **Example L2**

(2-Amino-4-iodo-5-morpholin-4-yl-phenyl)-carbamic acid tert.-butyl ester

Prepared from (4-iodo-5-morpholin-4-yl-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example C3) (753 g, 1.65 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (1.9 g, 8.27 mmol) according to the general procedure L (method b). Obtained as a yellow solid (696 mg).

MS (ISP) 420 $[(M+H)^+]$; mp 139-143 °C.

Example L3

(2-Amino-5-chloro-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester

- Prepared from (5-chloro-2-nitro-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester
5 (Example K1) (742 mg, 2.0 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (2.245 g, 10 mmol) according to the general procedure L (method b). Obtained as an orange solid (483 mg).

MS (ISP) 343 $[(M+H)^+]$ and 345 $[(M+2+H)^+]$.

Example L4

(2-Amino-5-methyl-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester

- 10 Prepared from (5-methyl-2-nitro-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example K2) (741 mg, 2.1 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (2.37 g, 10.3 mmol) according to the general procedure L (method b). Obtained as a light-brown solid (419 mg).

MS (ISP) 323 $[(M+H)^+]$; mp 172-173 °C.

15 Example L5

[2-Amino-5-(4-methyl-piperazin-1-yl)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester

- Prepared from [5-(4-methyl-piperazin-1-yl)-2-nitro-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example K3) (1.08 g, 2.48 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (2.8
20 g, 12.4 mmol) according to the general procedure L (method b). Obtained as an orange solid (1.0 g).

MS (ISP) 407 $[(M+H)^+]$; mp 81-85 °C.

Example L6

(5-Amino-2-chloro-biphenyl-4-yl)-carbamic acid tert.-butyl ester

- 25 Prepared from (2-chloro-5-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example F1) (783 mg, 2.24 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (2.53 g, 11.2 mmol) according to the general procedure L (method b). Obtained as a yellow solid (684 mg).

MS (EI) 318 (M^+) and 320 [$(M+2)^+$]; mp 109-111 °C.

Example L7

(5-Amino-2-methyl-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (2-methyl-5-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example
5 F2) (921 mg, 2.8 mmol) by catalytic hydrogenation with Pd/C according to the general
procedure L (method a). Obtained as a white solid (796 mg).

MS (EI) 298 (M^+); mp 122 °C.

Example L8

[2-Amino-5-(2-dimethylamino-ethylsulfanyl)-4-phenylethynyl-phenyl]-carbamic acid
10 tert.-butyl ester

Prepared from [5-(2-dimethylamino-ethylsulfanyl)-2-nitro-4-phenylethynyl-phenyl]-
carbamic acid tert.-butyl ester (Example K8) (551 mg, 1.25 mmol) by reduction with
SnCl₂·2H₂O (1.41 g, 6.25 mmol) according to the general procedure L (method b).
Obtained as an orange foam (510 mg).

15 MS (ISP) 412[($M+H$)⁺]; mp 115-117 °C.

Example L9

(4-Amino-5-tert.-butoxycarbonylamino-2-phenylethynyl-phenylsulfanyl)-acetic acid
methyl ester

Prepared from (5-tert.-butoxycarbonylamino-4-nitro-2-phenylethynyl-phenylsulfanyl)-
20 acetic acid methyl ester (Example K9) (634 mg, 1.43 mmol) by reduction with SnCl₂·2H₂O
(1.62 g, 7.16 mmol) according to the general procedure L (method b). Obtained as an
orange gum (590 mg).

MS (ISP) 413 [($M+H$)⁺], 435 [($M+Na$)⁺] and 451 [($M+K$)⁺].

Example L10

25 (4-Amino-5-tert.-butoxycarbonylamino-2-phenylethynyl-phenyl)-acetic acid methyl ester

Prepared from (5-tert.-butoxycarbonylamino-4-nitro-2-phenylethynyl-phenyl)-acetic acid
methyl ester (Example K16) (1.09 g, 2.66 mmol) by reduction with SnCl₂·2H₂O (3.00 g,

13.28 mmol) according to the general procedure L (method b). Obtained as a light yellow solid (900 mg).

MS (ISP) 381 [(M+H)⁺] and 403 [(M+Na)⁺]; mp 130 °C.

Example L11

- 5 [2-Amino-5-(2-methoxy-ethoxy)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester

Prepared from [5-(2-methoxy-ethoxy)-2-nitro-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example K10) (1.417 g, 3.44 mmol) by reduction with SnCl₂·2H₂O (3.88 g, 17.2 mmol) according to the general procedure L (method b). Obtained as an off-white solid (1.04 g).

- 10 MS (EI) 382 (M⁺); mp 105-107 °C.

Example L12

(2-Amino-5-morpholin-4-yl-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester

- Prepared from (5-morpholin-4-yl-2-nitro-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example K4) (563 mg, 1.329 mmol) by reduction with SnCl₂·2H₂O (1.5 g, 6.65 mmol) according to the general procedure L (method b). Obtained as a red-brown solid (488 mg).
- 15

MS (ISP) 394 [(M+H)⁺]; mp 174-176 °C.

Example L13

(2-Amino-5-methoxy-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester

- 20 Prepared from (5-methoxy-2-nitro-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example K11) (1.00 g, 2.71 mmol) by reduction with SnCl₂·2H₂O (3.06 g, 13.57 mmol) according to the general procedure L (method b). Obtained as a yellow solid (870 mg).

MS (EI) 338 (M⁺); mp 158 °C.

- 25 Example L14

4-Amino-5-tert.-butoxycarbonylamino-2-phenylethynyl-benzoic acid methyl ester

Prepared from (5-tert.-butoxycarbonylamino-4-nitro-2-phenylethynyl-benzoic acid methyl ester (Example K15) (754 mg, 1.90 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (2.15 g, 9.51 mmol) according to the general procedure L (method b). Obtained as a pink solid (431 mg).

5 MS (EI) 366 (M^+); mp 164 °C.

Example L15

[2-Amino-5-(2-[2-[2-(2-methoxy-ethoxy)-ethoxy]-ethoxy]-ethoxy)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester

Prepared from [5-(2-[2-[2-(2-methoxy-ethoxy)-ethoxy]-ethoxy]-ethoxy)-2-nitro-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example K12) (3.0 g, 5.51 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (6.2 g, 27.54 mmol) according to the general procedure L (method b). Obtained as a brown oil (2.9 g).

MS (ISP) 515 [$(\text{M}+\text{H})^+$].

Example L16

15 (4-Amino-5-tert.-butoxycarbonylamino-2-phenylethynyl-phenoxy)-acetic acid tert.-butyl ester

Prepared from (5-tert.-butoxycarbonylamino-4-nitro-2-phenylethynyl-phenoxy)-acetic acid tert.-butyl ester (Example K13) (1.32 g, 2.82 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (3.18 g, 14.10 mmol) according to the general procedure L (method b). Obtained as an amorphous orange material (1.2 g).

MS (ISP) 439 [$(\text{M}+\text{H})^+$] and 461 [$(\text{M}+\text{Na})^+$].

Example L17

(2-Amino-5-cyanomethyl-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester

Prepared from (5-cyanomethyl-2-nitro-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example K17) (377 mg, 1.0 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (1.13 g, 5.0 mmol) according to the general procedure L (method b). Obtained as an orange solid (338 mg).

MS (ISP) 348 [$(\text{M}+\text{H})^+$] and 370 [$(\text{M}+\text{Na})^+$]; mp 143 °C.

Example L18

[2-Amino-5-(1,4-dioxo-8-aza-spiro[4.5]dec-8-yl)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester and [2-Amino-5-(4,4-diethoxy-piperidin-1-yl)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester

- 5 Prepared from [5-(1,4-dioxo-8-aza-spiro[4.5]dec-8-yl)-2-nitro-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example K7) (1.73 g, 3.6 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (4.0 g, 18.0 mmol) in EtOH according to the general procedure L (method b). Obtained as a brown solid (418 mg) and as dark brown solid (379 mg), respectively.

MS (ISP) 450 [(M+H)⁺]; mp 79-82 °C; MS (ISP) 480 [(M+H)⁺].

10 Example L19

[2-Amino-5-(1,1-dioxo-6-thiomorpholin-4-yl)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester

- Prepared from [5-(1,1-dioxo-thiomorpholin-4-yl)-2-nitro-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example K6) (235 mg, 0.5 mmol) by reduction with
15 $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (564 mg, 2.5 mmol) according to the general procedure L (method b). Obtained as a brown solid (418 mg, impure material, used directly in Example 6 without purification and characterization).

Example L20

- [2-Amino-5-(2,2-dimethyl-[1,3]dioxolan-4-ylmethoxy)-4-phenylethynyl-phenyl]-
20 carbamic acid tert.-butyl ester

- Prepared from (RS)-[5-(2,2-dimethyl-[1,3]dioxolan-4-ylmethoxy)-2-nitro-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example B8) (1.90 g, 4.06 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (4.6 g, 20.32 mmol) according to the general procedure L (method b). The crude product was reprotected by stirring with 2,2-dimethoxypropane (5
25 mL) and p-TsOH·H₂O (1.1 eq) in DMF (5 mL) at 23 °C for 4 h. Obtained as a brown solid (1.1 g).

MS (ISP) 439 [(M+H)⁺], 461 [(M+Na)⁺] and 477 [(M+K)⁺].

Example L21

4-Amino-5-tert.-butoxycarbonylamino-2-iodo-benzoic acid methyl ester

Prepared from 5-tert.-butoxycarbonylamino-2-iodo-4-nitro-benzoic acid methyl ester (Example B3) (3.00 g, 7.11 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (8.02 g, 35.55 mmol) according to the general procedure L (method b). Obtained as a light red foam (1.9 g).

MS (ISP) 393 [(M+H)⁺]; mp 60-78 °C.

5 Example L22

[RS]-[2-Amino-5-(2-oxo-[1,3]dioxolan-4-ylmethoxy)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester

Prepared from [RS]-[2-Nitro-5-(2-oxo-[1,3]dioxolan-4-ylmethoxy)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester [RO-68-8108/000 (411 mg, 0.9 mmol)], prepared
10 from (RS)-[5-(2,3-dihydroxy-propoxy)-2-nitro-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example K18) by treatment with 1,1'-carbonyldiimidazole in THF at 0 to 23 °C] by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (1.02 g, 4.5 mmol) according to the general procedure L (method b). Obtained as an apricot solid (370 mg).

MS (ISP) 425 [(M+H)⁺]; mp 140 °C.

15 Example L23

(2-Amino-5-ethoxymethyl-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester

Prepared from carbonic acid 5-tert.-butoxycarbonylamino-4-nitro-2-phenylethynyl-benzyl ester methyl ester [RO-68-8481/000, prepared from (5-hydroxymethyl-2-nitro-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example K19) by treatment with
20 methyl chloroformate and Et_3N in THF at 0 °C] by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ according to the general procedure L (method b). Obtained as an amorphous brown substance (139 mg).

MS (ISP) 367 [(M+H)⁺].

Example L24

25 2,2-Dimethyl-propionic acid 4-amino-5-tert.-butoxycarbonylamino-2-phenylethynyl-benzyl ester

Prepared from 2,2-dimethyl-propionic acid 5-tert.-butoxycarbonylamino-4-nitro-2-phenylethynyl-benzyl ester [RO-68-9779/000, prepared from (5-hydroxymethyl-2-nitro-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example K19) by treatment with

pivaloyl chloride and cat. DMAP in pyridine at 0 to 23 °C] by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ according to the general procedure L (method b). Obtained as an amorphous yellow substance (182 mg).

MS (ISN) 421 [(M-H)⁻].

5 Example L25

(RS)-[2-Amino-4-phenylethynyl-5-(tetrahydro-pyran-2-yloxymethyl)-phenyl]-carbamic acid tert.-butyl ester

Prepared from (RS)-[2-nitro-4-phenylethynyl-5-(tetrahydro-pyran-2-yloxymethyl)-phenyl]-carbamic acid tert.-butyl ester [RO-69-2829/000, prepared from (5-hydroxymethyl-2-nitro-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example K19) by treatment with 3,4-dihydro-2H-pyran and cat. $\text{TsOH} \cdot \text{H}_2\text{O}$ in DCM at 0 °C] by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ according to the general procedure L (method b). Obtained as a yellow solid (1.78 g).

MS (ISN) 421 [(M-H)⁻]; mp 158 °C.

15 Example L26

[2-Amino-5-(4-methoxy-piperidin-1-yl)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester

Prepared from [5-(4-methoxy-piperidin-1-yl)-2-nitro-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example K20) (727 mg, 1.61 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ according to the general procedure L (method b). Obtained as a yellow solid (489 mg).

MS (ISP) 422 [(M+H)⁺]; mp 173-176 °C.

Example L27

(2-Amino-5-cyanomethoxy-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester

Prepared from (5-cyanomethoxy-2-nitro-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example K21) (395 mg, 0.91 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ according to the general procedure L (method b). Obtained as an amorphous yellow substance (219 mg).

MS (ISP) 364 [(M+H)⁺].

Example L28

(RS)-[2-Amino-4-(4-fluoro-phenylethynyl)-5-(tetrahydro-pyran-2-yloxymethyl)-phenyl]-carbamic acid tert.-butyl ester

- Prepared from (RS)-[4-(4-fluoro-phenylethynyl)-2-nitro-5-(tetrahydro-pyran-2-yloxymethyl)-phenyl]-carbamic acid tert.-butyl ester [RO-69-3877/000, prepared from [4-(4-fluoro-phenylethynyl)-5-hydroxymethyl-2-nitro-phenyl]-carbamic acid tert.-butyl ester (Example K22) by treatment with 3,4-dihydro-2H-pyran and cat. TsOH.H₂O in DCM at 0 °C] by reduction with SnCl₂·2H₂O according to the general procedure L (method b).
Obtained as an amorphous light brown substance (990 mg).

- 10 MS (ISP) 441 [(M+H)⁺].

Example L29

(RS)-(2-Amino-4-(4-fluoro-phenylethynyl)-5-{methyl-[2-(tetrahydro-pyran-2-yloxy)-ethyl]-amino}-phenyl)-carbamic acid tert.-butyl ester

- Prepared from (RS)-(4-(4-fluoro-phenylethynyl)-5-{methyl-[2-(tetrahydro-pyran-2-yloxy)-ethyl]-amino}-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example K23) (1.79 g, 3.49 mmol) by reduction with SnCl₂·2H₂O according to the general procedure L (method b). Obtained as an amorphous light brown substance (1.20 g).

MS (ISP) 484 [(M+H)⁺].

Example L30

- 20 (RS)-[2-Amino-4-phenylethynyl-5-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-phenyl]-carbamic acid tert.-butyl ester

- Prepared from (RS)-[2-nitro-4-phenylethynyl-5-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-phenyl]-carbamic acid tert.-butyl ester (Example K24) (420 mg, 0.87 mmol) by reduction with SnCl₂·2H₂O according to the general procedure L (method b). Obtained as a light brown solid (346 mg).

25 MS (ISP) 453 [(M+H)⁺].

Example L31

(RS)-{5-Amino-4'-fluoro-2-[4-(tetrahydro-pyran-2-yloxy)-piperidin-1-yl]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- Prepared from (RS)-{4'-fluoro-5-nitro-2-[4-(tetrahydro-pyran-2-yloxy)-piperidin-1-yl]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example F3) (845 mg, 1.64 mmol) by
5 catalytic hydrogenation with Pd/C according to the general procedure L (method a).
Obtained as a light green solid (758 mg).

MS (ISP) 486 [(M+H)⁺]; mp 157-161 °C.

Example L32

- 10 [2-Amino-5-(2-tert.-butoxy-ethoxy)-4-(4-fluoro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester

- Prepared from [5-(2-tert.-butoxy-ethoxy)-4-(4-fluoro-phenylethynyl)-2-nitro-phenyl]-carbamic acid tert.-butyl ester (Example K25) (744 mg, 1.57 mmol) by reduction with
SnCl₂·2H₂O according to the general procedure L (method b). Obtained as a light yellow
15 solid (575 mg).

MS (ISP) 443 [(M+H)⁺]; mp 149-150 °C.

Example L33

(RS)-{5-Amino-4'-fluoro-2-[4-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-piperidin-1-yl]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- 20 Prepared from (RS)-{4'-fluoro-5-nitro-2-[4-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-piperidin-1-yl]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example F4) (900 mg, 1.61 mmol) by catalytic hydrogenation with Pd/C according to the general procedure L (method a). Obtained as a light brown foam (779 mg).

MS (ISP) 530 [(M+H)⁺]; mp 56-58 °C.

- 25 Example L34

(RS)-{5-Amino-4'-fluoro-2-(tetrahydro-pyran-2-yloxymethyl)-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (RS)-[4'-fluoro-5-nitro-2-(tetrahydro-pyran-2-yloxymethyl)-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example F5) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ according to the general procedure L (method b). Obtained as an orange solid (1.15 g).

mp 139-142 °C.

5 Example L35

(5-Amino-2-cyanomethoxy-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (2-cyanomethoxy-4'-fluoro-5-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example F6) (310 mg, 0.8 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ according to the general procedure L (method b). Obtained as a light brown solid (220 mg).

10 MS (ISN) 356 [(M-H)⁻]; mp 118-119 °C.

Example L36

(5-Amino-2-dimethylaminomethyl-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (2-dimethylaminomethyl-4'-fluoro-5-nitro-biphenyl-4-yl)-carbamic acid
15 tert.-butyl ester (Example F7) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ according to the general procedure L (method b). Obtained as a yellow solid (908 mg).

mp 97-125 °C.

Example L37

[5-Amino-2-(2,2-dimethyl-tetrahydro-[1,3]dioxolo[4,5-c]pyrrol-5-yl)-4'-fluoro-biphenyl-
20 4-yl]-carbamic acid tert.-butyl ester

Prepared from [2-(2,2-dimethyl-tetrahydro-[1,3]dioxolo[4,5-c]pyrrol-5-yl)-4'-fluoro-5-nitro-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example F8) (610 mg, 1.29 mmol) by catalytic hydrogenation with Pd/C according to the general procedure L (method a). Obtained as an off-white foam (578 mg).

25 MS (ISP) 444 [(M+H)⁺].

Example L38

(5-Amino-4'-fluoro-2-methoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (4'-fluoro-2-methoxy-5-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example F9) (2.64 g, 7.29 mmol) by catalytic hydrogenation with Pd/C according to the general procedure L (method a). Obtained as an off-white solid (2.36 g).

MS (ISP) 333 [(M+H)⁺]; mp 155 °C (dec.).

5 Example L39

[5-Amino-2-(1,4-dioxo-8-aza-spiro[4.5]dec-8-yl)-4'-fluoro-biphenyl-4-yl]-carbamic acid tert.-butyl ester

Prepared from [2-(1,4-dioxo-8-aza-spiro[4.5]dec-8-yl)-4'-fluoro-5-nitro-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example F10) (2.4 g, 5.0 mmol) by catalytic hydrogenation
10 with Pd/C according to the general procedure L (method a). Obtained as a green solid (2.37 g).

MS (ISP) 444 [(M+H)⁺].

Example L40

(5-Amino-4'-fluoro-2-methyl-biphenyl-4-yl)-carbamic acid tert.-butyl ester

15 Prepared from (4'-fluoro-2-methyl-5-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example F11) (560 mg, 1.62 mmol) by catalytic hydrogenation with Pd/C according to the general procedure L (method a). Obtained as a light brown solid (512 mg).

MS (ISP) 317 [(M+H)⁺]; mp 112 °C.

Example L41

20 (5-Amino-4-tert.-butoxycarbonylamino-4'-fluoro-biphenyl-2-yloxy)-acetic acid tert.-butyl ester

Prepared from (4-tert.-butoxycarbonylamino-4'-fluoro-5-nitro-biphenyl-2-yloxy)-acetic acid tert.-butyl ester (Example F12) (2.29 g, 4.95 mmol) by catalytic hydrogenation with
25 Pd/C according to the general procedure L (method a). Obtained as a dark blue solid (2.14 g).

MS (ISP) 433 [(M+H)⁺]; mp 30-33 °C.

Example L42

(5-Amino-2-chloro-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (2-chloro-4'-fluoro-5-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example F13) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ according to the general procedure L

5 (method b). Obtained as a light red solid (544 mg).

MS (ISP) 337 $[(\text{M}+\text{H})^+]$.

Example L43

[5-Amino-4'-fluoro-2-(2-methoxy-ethoxy)-biphenyl-4-yl]-carbamic acid tert.-butyl ester

Prepared from [4'-fluoro-2-(2-methoxy-ethoxy)-5-nitro-biphenyl-4-yl]-carbamic acid
10 tert.-butyl ester (Example F14) by catalytic hydrogenation with Pd/C according to the
general procedure L (method a). Obtained as a light brown solid (1.652 g).

MS (ISP) 377 $[(\text{M}+\text{H})^+]$.

Example L44

[5-Amino-2-(2-tert.-butoxy-ethoxy)-4'-fluoro-biphenyl-4-yl]-carbamic acid tert.-butyl
15 ester

Prepared from [2-(2-tert.-butoxy-ethoxy)-4'-fluoro-5-nitro-biphenyl-4-yl]-carbamic acid
tert.-butyl ester (Example F15) by catalytic hydrogenation with Pd/C according to the
general procedure L (method a). Obtained as a purple solid (547 mg).

MS (ISP) 419 $[(\text{M}+\text{H})^+]$; mp 133 °C (dec.).

20 Example L45

[5-Amino-4'-fluoro-2-(2-oxo-oxazolidin-3-yl)-biphenyl-4-yl]-carbamic acid tert.-butyl
ester

Prepared from [4'-fluoro-5-nitro-2-(2-oxo-oxazolidin-3-yl)-biphenyl-4-yl]-carbamic acid
tert.-butyl ester (Example F16) (280 mg, 0.67 mmol) by catalytic hydrogenation with Pd/C
25 according to the general procedure L (method a). Obtained as a yellow solid (277 mg).

MS (ISP) 388 $[(\text{M}+\text{H})^+]$; mp 210 °C.

Example L46

(5-Amino-4'-fluoro-2-methoxy-2'-methyl-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (4'-fluoro-2-methoxy-2'-methyl-5-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example F17) by catalytic hydrogenation with Pd/C according to the general
5 procedure L (method a). Obtained as a brown solid (588 mg).

MS (ISP) 347 [(M+H)⁺].

Example L47

(5-Amino-2-tert.-butoxy-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (2-tert.-butoxy-4'-fluoro-5-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl
10 ester (Example F18) (1.15 g, 2.84 mmol) by catalytic hydrogenation with Pd/C according to the general procedure L (method a). Obtained as a pink solid (747 mg).

MS (ISP) 375 [(M+H)⁺]; mp 139 °C.

Example L48

(5-Amino-2-tert.-butoxy-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester

15 Prepared from (2-tert.-butoxy-2'-fluoro-5-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example F19) (930 mg, 2.3 mmol) by catalytic hydrogenation with Pd/C according to the general procedure L (method a). Obtained as a pink solid (649 mg).

MS (ISP) 375 [(M+H)⁺]; mp 130 °C.

Example L49

20 (RS)-{5-Amino-4'-fluoro-2-[(R)-3-(tetrahydro-pyran-2-yloxy)-pyrrolidin-1-yl]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (RS)-{4'-fluoro-5-nitro-2-[(R)-3-(tetrahydro-pyran-2-yloxy)-pyrrolidin-1-yl]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example F20) by catalytic
hydrogenation with Pd/C according to the general procedure L (method a). Obtained as a
25 dark green solid (852 mg).

MS (ISP) 472 [(M+H)⁺].

Example L50

(5-Amino-2'-fluoro-2-methoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester

- Prepared from (2'-fluoro-2-methoxy-5-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example F21) (649 mg, 1.79 mmol) by catalytic hydrogenation with Pd/C according to the general procedure L (method a). Obtained as a light brown solid (352 mg).

MS (ISP) 333 [(M+H)⁺]; mp 161 °C.

Example L51

[5-Amino-2-(1,4-dioxo-8-aza-spiro[4.5]dec-8-yl)-2'-fluoro-biphenyl-4-yl]-carbamic acid tert.-butyl ester

- 10 Prepared from [2-(1,4-dioxo-8-aza-spiro[4.5]dec-8-yl)-2'-fluoro-5-nitro-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example F22) (1.39 g, 2.94 mmol) by catalytic hydrogenation with Pd/C according to the general procedure L (method a). Obtained as a light beige solid (1.01 g).

MS (ISP) 444 [(M+H)⁺]; mp 198 °C.

15 Example L52

(5-Amino-2',5'-difluoro-2-methoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (2',5'-difluoro-2-methoxy-5-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example F23) (1.05 g, 2.76 mmol) by catalytic hydrogenation with Pd/C according to the general procedure L (method a). Obtained as a beige solid (618 mg).

- 20 MS (ISN) 349 [(M-H)⁻]; mp 144 °C.

Example L53

[5-Amino-2'-fluoro-2-(2-methoxy-ethoxy)-biphenyl-4-yl]-carbamic acid tert.-butyl ester

- Prepared from [2'-fluoro-2-(2-methoxy-ethoxy)-5-nitro-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example F24) by catalytic hydrogenation with Pd/C according to the general procedure L (method a). Obtained as a purple solid (2.581 g).

Example L54

(RS)-[5-Amino-2'-fluoro-2-(tetrahydro-pyran-2-yloxymethyl)-biphenyl-4-yl]-carbamic acid tert.-butyl ester

Prepared from (RS)-[2'-fluoro-5-nitro-2-(tetrahydro-pyran-2-yloxymethyl)-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example F25) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ according to the general procedure L (method b). Obtained as a yellow liquid (2.676 g).

MS (ISP) 439 $[(\text{M}+\text{Na})^+]$.

Example L55

[5-Amino-2'-fluoro-2-(4-methoxy-benzyloxy)-biphenyl-4-yl]-carbamic acid tert.-butyl ester

Prepared from [2'-fluoro-2-(4-methoxy-benzyloxy)-5-nitro-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example F25) (0.90 g, 1.92 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ according to the general procedure L (method b). Obtained as a beige solid (719 mg).

MS (ISP) 439 $[(\text{M}+\text{H})^+]$.

Example L56

(5-Amino-2-tert.-butoxy-2',5'-difluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared from (2-tert.-butoxy-2',5'-difluoro-5-nitro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example F27) by catalytic hydrogenation with Pd/C according to the general procedure L (method a). Obtained as an amorphous grey-blue substance (2.37 g).

MS (ISP) 393 $[(\text{M}+\text{H})^+]$.

Example L57

[2-Amino-5-tert.-butoxy-4-(4-fluoro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester

Prepared from [5-tert.-butoxy-4-(4-fluoro-phenylethynyl)-2-nitro-phenyl]-carbamic acid tert.-butyl ester (Example K26) (649 mg, 1.51 mmol) by reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ according to the general procedure L (method b). Obtained as a light yellow solid (410 mg).

MS (ISP) 399 [(M+H)⁺]; mp 183 °C.

The following examples relate to the preparation of the ethyl or tert.-butyl 3-aryl-3-oxo-propionates (general formula VIIa), which serve as building blocks in the synthesis of the target compounds (Synthetic Scheme H):

5 General procedure M

Method a) Preparation of ethyl or tert.-butyl 3-aryl-3-oxo-propionates

The ethyl or tert.-butyl 3-aryl-3-oxo-propionates were prepared from the aryl acid chlorides and ethyl or tert.-butyl malonate potassium salt [CAS-no. 6148-64-7 and 75486-33-8] with Et₃N and MgCl₂ in CH₃CN at 0 °C to 23 °C according to *Synthesis* 1993, 290. If
10 the free aryl carboxylic acid was employed in this reaction, it was activated by treatment with ethyl chloroformate and Et₃N in THF/CH₃CN at 0 °C prior to reaction with the malonate salt.

Method b) Preparation of tert.-butyl 3-aryl-3-oxo-propionates

The tert.-butyl 3-aryl-3-oxo-propionates were alternatively prepared from the methyl or
15 ethyl aryl esters by treatment with lithium tert.-butyl acetate [prepared by treatment of tert.-butyl acetate with lithium diisopropylamide in THF at -78 °C] in the presence of lithium tert.-butoxide according to *Synthesis* 1985, 45. If the product contained residual starting material after workup, thus could be removed by selective saponification with LiOH in THF/MeOH/H₂O at 23 °C.


20 Method c) Preparation of 3-aryl-3-oxo-propionic acids

The 3-aryl-3-oxo-propionic acids were prepared from the aryl acid chlorides and bis(trimethylsilyl)malonate with Et₃N and LiBr in CH₃CN at 0 °C according to *Synth. Commun.* 1985, 15, 1039 (method c1) or with n-BuLi in ether at -60 °C to 0 °C according to *Synthesis* 1979, 787 (method c2).

25 Example M1

3-Oxo-3-(3-[1,2,4]triazol-4-yl-phenyl)-propionic acid ethyl ester.

RO-71-2790/000



- Prepared from 3-[1,2,4]triazol-4-yl-benzoic acid [RO-71-1432/000, prepared by reaction of 3-aminobenzoic acid with hydrazine hydrate and triethyl orthoformate in acetic acid at 120 °C] by activation with ethyl chloroformate/Et₃N and reaction with ethyl malonate potassium salt with Et₃N and MgCl₂ in CH₃CN according to general procedure M (method a). Obtained as a white solid (5.74 g).

MS (EI) 259 (M⁺).Example M2

- 3-Oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionic acid ethyl ester.

Prepared from 3-[1,2,3]triazol-1-yl-benzoic acid [RO-71-3703/000, prepared by refluxing of methyl 3-azidobenzoate [CAS-No. 93066-93-4] in trimethylsilylacetylene, followed by saponification with aqueous NaOH in refluxing EtOH] by activation with ethyl chloroformate/Et₃N and reaction with ethyl malonate potassium salt with Et₃N and MgCl₂ in CH₃CN according to general procedure M (method a). Obtained as a light yellow solid (2.22 g).

MS (EI) 259 (M⁺); mp 72-74 °C.Example M3

3-(3-Cyano-phenyl)-3-oxo-propionic acid tert.-butyl ester.

- Prepared from methyl 3-cyanobenzoate [CAS-No. 13531-48-1] by treatment with lithium tert.-butyl acetate according to general procedure M (method b). Obtained as a light brown oily semisolid.

MS (EI) 245 (M⁺).Example M4

- 3-(3-Imidazol-1-yl-phenyl)-3-oxo-propionic acid tert.-butyl ester.

Prepared from methyl 3-(1H-imidazol-1-yl)benzoate [prepared from 3-(1H-imidazol-1-yl)benzoic acid (*J. Med. Chem.* 1987, 30, 1342; CAS-No. [108035-47-8] by refluxing in conc. H₂SO₄/MeOH] by treatment with lithium tert.-butyl acetate according to general procedure M (method b). Obtained as an orange-brown oil.

5 MS (ISP) 287 [(M+H)⁺].

Example M5

3-(2-Imidazol-1-yl-pyridin-4-yl)-3-oxo-propionic acid tert.-butyl ester.

Prepared from 2-imidazol-1-yl-isonicotinoyl chloride hydrochloride [prepared by reaction of tert.-butyl 2-chloroisonicotinoate with imidazole and NaH in DMF at 80 °C, treatment
10 with formic acid at 50 °C and reaction with thionylchloride in toluene at 100 °C] and tert.-butyl malonate potassium salt with Et₃N and MgCl₂ in CH₃CN according to general procedure M (method a). Obtained as a brown solid (10.8 g).

MS (EI) 287 (M⁺); mp 80 °C (dec.).

Example M6

15 3-Oxo-3-(3-[1,2,4]triazol-1-yl-phenyl)-propionic acid tert.-butyl ester.

Prepared from methyl 3-[1,2,4]triazol-1-yl-benzoate [CAS-No. 167626-27-9] by treatment with lithium tert.-butyl acetate according to general procedure M (method b). Obtained as an orange liquid (2.41 g).

MS (EI) 287 (M⁺).

20 Example M7

3-[3-(4-Methyl-imidazol-1-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester.

Prepared from methyl 3-(4-methyl-imidazol-1-yl)-benzoate [RO-69-6483/000, prepared the corresponding acid from 3-isothiocyanatobenzoic acid and 2-aminopropionaldehyde dimethyl acetal according to *J. Med. Chem.* 1987, 30, 1342, followed by refluxing in conc.
25 H₂SO₄/MeOH] by treatment with lithium tert.-butyl acetate according to general procedure M (method b). Obtained as a yellow-brown oil (10.69 g).

MS (EI) 300 (M⁺).

Example M8

3-[3-(2-Methyl-imidazol-1-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester.

- Prepared from ethyl 3-(2-methyl-imidazol-1-yl)-benzoate [RO-69-7480/000, prepared by reaction of ethyl 3-aminobenzoate with ethyl acetimidate hydrochloride in EtOH at 0 °C, direct treatment with aminoacetaldehyde diethyl acetal in EtOH at 23 °C, followed by addition of conc. H₂SO₄ and refluxing.] by treatment with lithium tert.-butyl acetate according to general procedure M (method b). Obtained as a brown oil (9.66 g).

MS (ISN) 299 [(M-H)⁺].

Example M9

- 10 3-[3-(2,4-Dimethyl-imidazol-1-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester.

- Prepared from ethyl 3-(2,4-dimethyl-imidazol-1-yl)-benzoate [RO-71-0583/000, prepared by reaction of ethyl 3-aminobenzoate with ethyl acetimidate hydrochloride in EtOH at 0 °C, direct treatment with 2-aminopropionaldehyde dimethyl acetal in EtOH at 23 °C, followed by addition of conc. H₂SO₄ and refluxing.] by treatment with lithium tert.-butyl acetate according to general procedure M (method b). Obtained as a yellow-brown oil (6.00 g).

MS (ISN) 313 [(M-H)⁺].

Example M10

3-(2-Cyano-pyridin-4-yl)-3-oxo-propionic acid tert.-butyl ester.

- 20 Prepared from 2-cyano-isonicotinic acid ethyl ester [CAS-No. 58481-14-4] by treatment with lithium tert.-butyl acetate according to general procedure M (method b). Obtained as a light brown solid (7.70 g).

MS (ISN) 245 [(M-H)⁺].

Example M11

- 25 3-Oxo-3-(3-[1,2,4]triazol-4-yl-phenyl)-propionic acid tert.-butyl ester.

Prepared from methyl 3-[1,2,4]triazol-4-yl-benzoate [prepared by reaction of 3-aminobenzoic acid with hydrazine hydrate and triethyl orthoformate in acetic acid at 120 °C, followed by esterification with conc. H₂SO₄ in refluxing MeOH] by treatment with

lithium tert.-butyl acetate according to general procedure M (method b). Obtained as a light yellow gum (870 mg).

MS (ISN) 286 [(M-H)⁻].

Example M12

- 5 3-[3-(2-Methoxymethylsulfanyl-imidazol-1-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester.

Prepared from ethyl 3-(2-methoxymethylsulfanyl-imidazol-1-yl)-benzoate [prepared by esterification of 3-(2-methoxymethylsulfanyl-imidazol-1-yl)-benzoic acid [CAS-No. 108035-46-7] with conc. H₂SO₄ in EtOH, followed by treatment with chloromethylmethyl
10 ether and NaH in THF/DMF] by treatment with lithium tert.-butyl acetate according to general procedure M (method b). Obtained as an orange oil (1.82 g).

MS (EI) 362 (M⁺).

Example M13

3-[3-(2-Methylsulfanyl-imidazol-1-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester.

- 15 Prepared from ethyl 3-(2-methylsulfanyl-imidazol-1-yl)-benzoate [prepared by esterification of 3-(2-methoxymethylsulfanyl-imidazol-1-yl)-benzoic acid [CAS-No. 108035-46-7] with conc. H₂SO₄ in EtOH, followed by treatment methyl iodide and NaH in THF/DMF] by treatment with lithium tert.-butyl acetate according to general procedure M (method b). Obtained as a light brown oil (4.41 g).

- 20 MS (ISP) 333 [(M+H)⁺].

Example M14

3-[3-(3-Methyl-isoxazol-5-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester.

- Prepared from ethyl 3-(3-methyl-isoxazol-5-yl)-benzoate [prepared by reaction of ethyl 3-ethynylbenzoate [CAS-No. 178742-95-5] with a mixture of NCS, acetaldoxime, Et₃N and
25 cat. amount of pyridine in CHCl₃ at 50 °C according to *Tetrahedron* 1984, 40, 2985-2988] by treatment with lithium tert.-butyl acetate according to general procedure M (method b). Obtained as a yellow solid (2.54 g).

MS (ISP) 302 [(M+H)⁺]; mp 50-56 °C.

Example M15

3-Oxo-3-(3-tetrazol-1-yl-phenyl)-propionic acid ethyl ester.

- Prepared from 3-tetrazol-1-yl-benzoic acid [CAS-No. 204196-80-5] by activation with ethyl chloroformate/ Et_3N and reaction with ethyl malonate potassium salt with Et_3N and MgCl_2 in CH_3CN according to general procedure M (method a). Obtained as a light yellow solid (211 mg).

MS (EI) 260 (M^+).

Example M16

3-(3-Chloro-thiophen-2-yl)-3-oxo-propionic acid ethyl ester

- 10 Prepared from 3-chloro-2-thiophenecarbonyl chloride [CAS-No. 86427-02-3] by reaction with ethyl malonate potassium salt with Et_3N and MgCl_2 in CH_3CN according to general procedure M (method a). Obtained as a brown oil (6.84 g).

MS (EI) 232 (M^+) and 234 [$(\text{M}+2)^+$].

Example M17

- 15 3-(5-Cyano-thiophen-2-yl)-3-oxo-propionic acid tert.-butyl ester

Prepared from ethyl 5-cyano-2-thiophenecarboxylate [CAS-No. 67808-35-9] by treatment with lithium tert.-butyl acetate according to general procedure M (method b). Obtained as a yellow solid (6.66 g).

MS (EI) 251 (M^+); mp 78 °C.

- 20 Example M18

3-(5-Cyano-2-fluoro-phenyl)-3-oxo-propionic acid ethyl ester

- Prepared from 5-cyano-2-fluoro-benzoyl chloride [prepared from the corresponding acid [CAS-No. 146328-87-2] by treatment with SOCl_2 , cat. DMF in toluene at 80 °C] by reaction with ethyl malonate potassium salt with Et_3N and MgCl_2 in CH_3CN according to general procedure M (method a). Obtained as a light yellow solid (3.85 g).

MS (EI) 235 (M^+); mp 55-60 °C.

Example M19

3-(2-Imidazol-1-yl-thiazol-4-yl)-3-oxo-propionic acid tert.-butyl ester

Prepared from ethyl 2-imidazol-1-yl-thiazole-4-carboxylate [CAS-No. 256420-32-3] by treatment with lithium tert.-butyl acetate according to general procedure M (method b).

- 5 Obtained as an orange oil (12.0 g).

Example M20

3-[2-(4-Methyl-imidazol-1-yl)-thiazol-4-yl]-3-oxo-propionic acid tert.-butyl ester

- Prepared from ethyl 2-(4-methyl-imidazol-1-yl)-thiazole-4-carboxylate [prepared from ethyl 2-amino-4-thiazolecarboxylate (CAS-No. [256420-32-3]) by the following synthetic sequence: 1.) NaH, 2-isothiocyanato-1,1-dimethoxy-propane, DMF, 23 °C; 2.) aq. H₂SO₄, reflux; 3.) EtOH, conc. H₂SO₄, 23 °C; 4.) 30% H₂O₂, HOAc, 23 °C] by treatment with lithium tert.-butyl acetate according to general procedure M (method b). Obtained as a brown oil (8.73 g).
- 10

MS (EI) 307 (M⁺).

15 Example M21

3-[3-(1-Methyl-1H-imidazol-2-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester

Prepared from ethyl 3-(1-methyl-1H-imidazol-2-yl)benzoate [CAS-No. 168422-44-4] by treatment with lithium tert.-butyl acetate according to general procedure M (method b). Obtained as a light yellow liquid (1.26 g).

- 20 MS (ISP) 301.3 [(M+H)⁺].

The following examples relate to the preparation of the 6-aryl-2,2-dimethyl-[1,3]dioxin-4-ones (general formula VII), which serve as building blocks in the synthesis of the target compounds (Synthetic Scheme H):

General procedure N

- 25 Preparation of 6-aryl-2,2-dimethyl-[1,3]dioxin-4-ones

Method a)

The 6-aryl-2,2-dimethyl-[1,3]dioxin-4-ones were prepared from 3-aryl-3-oxo-propionic acids and catalytic amount of conc. H_2SO_4 or trifluoroacetic acid (TFA) in isopropenyl acetate at 23 °C according to *Chem. Pharm. Bull.* 1983, 31, 1896. The final products were purified by silica gel column chromatography with hexane/EtOAc.

5 Method b)

The 6-aryl-2,2-dimethyl-[1,3]dioxin-4-ones were prepared from the tert.-butyl 3-aryl-3-oxo-propionates by treatment with trifluoroacetic anhydride (TFAA) in a mixture of TFA and acetone at 23 °C according to *Tetrahedron Lett.* 1998, 39, 2253. The final products were if necessary purified by silica gel column chromatography with hexane/EtOAc.

10 Example N1

2,2-Dimethyl-6-thiophen-2-yl-[1,3]dioxin-4-one.

The 3-oxo-3-thiophen-2-yl-propionic acid was prepared from thiophene-2-carbonyl chloride (5.3 mL, 50 mmol) and bis(trimethylsilyl)malonate (25.6 mL, 100 mmol) with n-BuLi (1.6M in hexane, 62.5 mL) in ether at -60 °C to 0 °C according to the general
15 procedure M (method c2). The crude material (7.88 g) was transformed into the title compound by stirring in isopropenyl acetate and TFA according to the general procedure N (method a). Obtained as a yellow solid (4.09 g).

MS (EI) 210 (M^+); mp 42 °C (dec.).

Example N2

20 6-(3-Chloro-thiophen-2-yl)-2,2-dimethyl-[1,3]dioxin-4-one.

The 3-(3-chloro-thiophen-2-yl)-3-oxo-propionic acid was prepared from 3-chloro-thiophene-2-carbonyl chloride (7.82 g, 43.2 mmol) and bis(trimethylsilyl)malonate (11.6 mL, 45.4 mmol) with Et_3N (12.65 mL, 90.7 mmol) and LiBr (3.53 g, 47.5 mmol) in CH_3CN at 0 °C according to general procedure M (method c1). The crude material (5.69 g) was
25 transformed into the title compound by stirring in isopropenyl acetate and conc. H_2SO_4 according to general procedure N (method a). Obtained as an orange solid (2.3 g).

MS (EI) 244 (M^+) and 246 [$(\text{M}+2)^+$]; mp 88-89 °C (dec.).

Example N3

6-(3-Cyano-thiophen-2-yl)-2,2-dimethyl-[1,3]dioxin-4-one.

- The 3-(3-cyano-thiophen-2-yl)-3-oxo-propionic acid was prepared from 3-cyano-thiophene-2-carbonyl chloride (24.33 g, 140.6 mmol) and bis(trimethylsilyl)malonate (38.0 mL, 147.7 mmol) with Et₃N (41 mL, 295.4 mmol) and LiBr (13.5 g, 154.7 mmol) in CH₃CN at 0 °C according to general procedure M (method c1). The crude material (24.8 g) was transformed into the title compound by stirring in isopropenyl acetate and conc. H₂SO₄ according to general procedure N (method a). Obtained as an orange solid (5.6 g).
- MS (EI) 235 (M⁺); mp 116-120 °C (dec.).

Example N4

3-(2,2-Dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile.

- 10 The 3-(3-cyano-phenyl)-3-oxo-propionic acid was prepared from 3-cyanobenzoyl chloride (828 mg, 5 mmol) and bis(trimethylsilyl)malonate (2.56 mL, 10 mmol) with n-BuLi (1.6M in hexane, 6.25 mL) in ether at -60 °C to 0 °C according to general procedure M (method c2). The crude material (1.04 g) was transformed into the title compound by stirring in isopropenyl acetate and TFA according to general procedure N (method a). Obtained as a
- 15 light yellow solid (0.8 g).
- MS (EI) 229 (M⁺); mp 138 °C (dec.).

Example N5

2,2-Dimethyl-6-(3-trifluoromethyl-phenyl)-[1,3]dioxin-4-one.

- 20 The 3-oxo-3-(3-trifluoromethyl-phenyl)-propionic acid was prepared from 3-trifluoromethylbenzoyl chloride (10 mL, 67.6 mmol) and bis(trimethylsilyl)malonate (18.2 mL, 71 mmol) with Et₃N (20 mL, 142 mmol) and LiBr (6.46 g, 74.4 mmol) in CH₃CN at 0 °C according to general procedure M (method c1). The crude material (7.0 g of the obtained 15.4 g) was transformed into the title compound by stirring in isopropenyl acetate and conc. H₂SO₄ according to general procedure N (method a). Obtained as a light
- 25 yellow solid (5.3 g).
- MS (EI) 272 (M⁺); mp 77-78 °C (dec.).

Example N6

6-(3-Chloro-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one.

The 3-(3-chloro-phenyl)-3-oxo-propionic acid was prepared from 3-chlorobenzoyl chloride (11 mL, 85.7 mmol) and bis(trimethylsilyl)malonate (23.0 mL, 90.0 mmol) with Et₃N (25 mL, 180 mmol) and LiBr (8.19 g, 94.3 mmol) in CH₃CN at 0 °C according to general procedure M (method c1). The crude material (17.1 g) was transformed into the title compound by stirring in isopropenyl acetate and conc. H₂SO₄ according to general procedure N (method a). Obtained as a yellow-brown solid (8.0 g).

MS (EI) 238 (M⁺) and 240 [(M+2)⁺]; mp 87-88 °C (dec.).

Example N7

6-(3-Iodo-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one.

10 The 3-(3-iodo-phenyl)-3-oxo-propionic acid was prepared from 3-iodobenzoyl chloride (21.0 g, 78.8 mmol) and bis(trimethylsilyl)malonate (21.0 mL, 82.8 mmol) with Et₃N (23 mL, 165.5 mmol) and LiBr (7.54 g, 86.7 mmol) in CH₃CN at 0 °C according to general procedure M (method c1). The crude material (21.9 g) was transformed into the title compound by stirring in isopropenyl acetate and conc. H₂SO₄ according to general procedure N (method a). Obtained as a yellow solid (9.6 g).

MS (EI) 330 (M⁺); mp 79-80 °C (dec.).

Example N8

2,2-Dimethyl-6-(3-trifluoromethoxy-phenyl)-[1,3]dioxin-4-one.

20 The 3-oxo-3-(3-trifluoromethoxy-phenyl)-propionic acid was prepared from 3-trifluoromethoxybenzoyl chloride and bis(trimethylsilyl)malonate with Et₃N and LiBr in CH₃CN at 0 °C according to general procedure M (method c1). The crude material was transformed into the title compound by stirring in isopropenyl acetate and conc. H₂SO₄ according to general procedure N (method a). Obtained as an orange solid (2.27 g).

MS (EI) 288 (M⁺); mp 49-54 °C (dec.).

25 Example N9

2,2-Dimethyl-6-pyridin-4-yl-[1,3]dioxin-4-one.

Prepared from 3-oxo-3-pyridin-4-yl-propionic acid [prepared from 4-acetylpyridine, magnesium methylcarbonate and CO₂ in DMF at 120 °C according to *Journal of Antibiotics*

1978, 31, 1245] by treatment with acetone, TFA and TFAA according to general procedure N (method b). Obtained as a white solid (1.3 g).

MS (EI) 205 (M^+)

Example N10

- 5 6-(3-Imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one.

The 3-(3-imidazol-1-yl-phenyl)-3-oxo-propionic acid was prepared from 3-(1H-imidazol-1-yl)benzoyl chloride hydrochloride [prepared by treatment of 3-(1H-imidazol-1-yl)benzoic acid (*J. Med. Chem.* 1987, 30, 1342; CAS-No. [108035-47-8] with SOCl_2) and bis(trimethylsilyl)malonate with Et_3N and LiBr in CH_3CN at 0 °C according to general
10 procedure M (method c1). The crude material was transformed into the title compound by stirring in isopropenyl acetate and conc. H_2SO_4 according to general procedure N (method a). Obtained as an orange semisolid (617 mg).

MS (EI) 270 (M^+).

Example N11

- 15 2,2-Dimethyl-6-(3-methoxy-phenyl)-[1,3]dioxin-4-one.

The 3-(3-methoxy-phenyl)-3-oxo-propionic acid was prepared from 3-methoxybenzoyl chloride (10.3 g, 60.4 mmol) and bis(trimethylsilyl)malonate (16.2 mL, 63.4 mmol) with Et_3N (17.7 mL, 127 mmol) and LiBr (5.77 g, 66.4 mmol) in CH_3CN at 0 °C according to
20 general procedure M (method c1). The crude material (6.38 g) was transformed into the title compound by stirring in isopropenyl acetate and conc. H_2SO_4 according to general procedure N (method a). Obtained as a yellow oil (640 mg).

MS (ISP) 235 [$(M+H)^+$] and 252 [$(M+\text{NH}_4)^+$].

Example N12

2,2-Dimethyl-6-(3-nitro-phenyl)-[1,3]dioxin-4-one.

- 25 The 3-(3-nitro-phenyl)-3-oxo-propionic acid tert.-butyl ester was prepared from 3-nitrobenzoyl chloride (2.71 g, 14.6 mmol) and tert.-butyl malonate potassium salt (6.0 g, 30.0 mmol) with Et_3N (4.5 mL, 32.2 mmol) and MgCl_2 (3.48 g, 36.52 mmol) in CH_3CN according to general procedure M (method a). The crude material (3.88 g) was

transformed into the title compound by stirring in TFA/acetone with TFAA according to general procedure N (method b). Obtained as a yellow solid (2.76 g).

MS (EI) 249 (M^+); mp 110-117 °C.

Example N13

- 5 2,2-Dimethyl-6-(3-[1,2,4]triazol-1-yl-phenyl)-[1,3]dioxin-4-one.

The 3-oxo-3-(3-[1,2,4]triazol-1-yl-phenyl)-propionic acid tert.-butyl ester [RO-69-3506/000] was prepared from 3-[1,2,4]triazol-1-yl-benzoic acid methyl ester [CAS-No. 167626-27-9] by treatment with lithium tert.-butyl acetate according to general procedure M (method b). Prepared from (Example M6) by stirring in TFA/acetone with TFAA
10 according to general procedure N (method b). Obtained as a yellow solid (539 mg).

MS (EI) 271 (M^+).

Example N14

- 6-(2-Imidazol-1-yl-pyridin-4-yl)-2,2-dimethyl-[1,3]dioxin-4-one.

Prepared from 3-(2-imidazol-1-yl-pyridin-4-yl)-3-oxo-propionic acid tert.-butyl ester
15 (Example M5) by stirring in TFA/acetone with TFAA according to general procedure N (method b). Obtained as a brown solid (10.8 g).

MS (EI) 271 (M^+); mp 151 °C (dec.).

Example N15

- 2,2-Dimethyl-6-[3-(2-methyl-imidazol-1-yl)-phenyl]-[1,3]dioxin-4-one.

20 Prepared from 3-[3-(2-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester (Example M8) by stirring in TFA/acetone with TFAA according to general procedure N (method b). Obtained as a beige solid (2.13 g).

MS (EI) 284 (M^+); mp 122 °C.

Example N16

- 25 4-(2,2-Dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-pyridine-2-carbonitrile.

Prepared from 3-(2-cyano-pyridin-4-yl)-3-oxo-propionic acid tert.-butyl ester (Example M10) by stirring in TFA/acetone with TFAA according to general procedure N (method b). Obtained as a brown solid (3.30 g).

MS (EI) 230 (M^+); mp 132 °C (dec.).

- 5 The following examples relate to the preparation of the 4,8-diaryl-1,3-dihydro-benzo[b][1,4]diazepin-2-ones, respectively the 4-aryl-8-arylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-ones and the 8-aryl-4-aryl-1,3-dihydro-benzo[b][1,4]diazepin-2-ones in regioisomerically pure fashion (Synthetic Scheme A):

General procedure O:

- 10 Preparation of {2-[3-aryl-3-oxo-propionylamino]-4-aryl-phenyl}-carbamic acid tert.-butyl ester by reaction of (2-amino-4-aryl-phenyl)-carbamic acid tert.-butyl esters with ethyl 3-aryl-3-oxo-propionates or 6-aryl-2,2-dimethyl-[1,3]dioxin-4-ones; also 3-aryl-N-(2-nitro-4-aryl-phenyl)-3-oxo-propionamides by reaction of 2-nitro-4-aryl-phenylamines with 6-aryl-2,2-dimethyl-[1,3]dioxin-4-ones
- 15 A mixture of the (2-amino-4-aryl-phenyl)-carbamic acid tert.-butyl ester or 2-nitro-4-aryl-phenylamine (1.0 mmol) and excess (1.2-1.5 mmol) of the ethyl 3-aryl-3-oxo-propionate [prepared from the aryl acid chloride and ethyl malonate potassium salt with Et₃N and MgCl₂ in CH₃CN at 23 °C according to *Synthesis* 1993, 290] or 6-aryl-2,2-dimethyl-[1,3]dioxin-4-one was refluxed in toluene (8 mL) until tlc indicated complete
- 20 consumption of the amine. The solution was allowed to cool to 23 °C, whereupon the product generally crystallized (in cases where crystallization failed to appear it was induced by addition of hexane). The solid was filtered off, washed with ether or mixtures of ether/hexane and dried in vacuum to give the {2-[3-aryl-3-oxo-propionylamino]-4-aryl-phenyl}-carbamic acid tert.-butyl esters or 3-aryl-N-(2-nitro-4-aryl-phenyl)-3-oxo-
- 25 propionamides, which was used directly in the following step or - if necessary - was purified by recrystallization or by silica gel column chromatography.

Example O1

{2-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-4-iodo-5-thiomorpholin-4-yl-phenyl}-carbamic acid tert.-butyl ester

- 30 Prepared from (2-amino-4-iodo-5-thiomorpholin-4-yl-phenyl)-carbamic acid tert.-butyl ester (Example L1) (653 mg, 1.5 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (690 mg, 2.25 mmol) according to the general procedure O. Obtained as a yellow solid (629 mg).

MS (ISP) 607 [(M+H)⁺].

Example O2

{2-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-4-iodo-5-morpholin-4-yl-phenyl}-carbamic acid tert.-butyl ester

- 5 Prepared from (2-amino-4-iodo-5-morpholin-4-yl-phenyl)-carbamic acid tert.-butyl ester (Example L2) (690 mg, 1.65 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (566 mg, 2.47 mmol) according to the general procedure O. Obtained as an orange solid (523 mg).

MS (ISP) 591 [(M+H)⁺] and 613 [(M+Na)⁺].

10 Example O3

{2-Chloro-5-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- Prepared from (5-amino-2-chloro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L6) (652 mg, 2.05 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (564 mg, 2.46 mmol) according to the general procedure O. Obtained as an off-white solid (725 mg).

MS (ISN) 488 [(M-H)⁻].

Example O4

- 20 {2-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-5-(2-dimethylamino-ethylsulfanyl)-4-phenylethynyl-phenyl}-carbamic acid tert.-butyl ester

Prepared from [2-amino-5-(2-dimethylamino-ethylsulfanyl)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example L8) (206 mg, 0.5 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (138 mg, 0.6 mmol) according to the general procedure O. Obtained as a yellow solid (210 mg).

- 25 MS (ISP) 583 [(M+H)⁺]; mp 88 °C.

Example O5

{5-tert.-Butoxycarbonylamino-4-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-2-phenylethynyl-phenylsulfanyl}-acetic acid methyl ester

Prepared from (4-amino-5-tert.-butoxycarbonylamino-2-phenylethynyl-phenylsulfanyl)-acetic acid methyl ester (Example L9) (534 mg, 1.3 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (358 mg, 1.56 mmol) according to the general procedure O. Obtained as a yellow foam (457 mg).

- 5 MS (ISP) 584 [(M+H)⁺], 601 [(M+NH₄)⁺] and 605 [(M+Na)⁺]; mp 69-73 °C.

Example O6

{5-tert.-Butoxycarbonylamino-4-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-2-phenylethynyl-phenyl}-acetic acid methyl ester

- 10 Prepared from (4-amino-5-tert.-butoxycarbonylamino-2-phenylethynyl-phenyl)-acetic acid methyl ester (Example L10) (721 mg, 2.0 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (550 mg, 2.4 mmol) according to the general procedure O. Obtained as a light yellow solid (886 mg).

MS (ISP) 552 [(M+H)⁺] and 569 [(M+NH₄)⁺]; mp 138 °C.

Example O7

- 15 [2-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-5-(2-methoxy-ethoxy)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester

- Prepared from [2-amino-5-(2-methoxy-ethoxy)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example L11) (415 mg, 1.09 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (373 mg, 1.63 mmol) according to the general
20 procedure O. Obtained as a light yellow solid (137 mg).

MS (ISP) 554 [(M+H)⁺], 571 [(M+NH₄)⁺] and 576 [(M+Na)⁺]; mp 175-176 °C.

Example O8

{2-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-5-methoxy-4-phenylethynyl-phenyl}-carbamic acid tert.-butyl ester

- 25 Prepared from (2-amino-5-methoxy-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example L12) (338 mg, 1.0 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (252 mg, 1.1 mmol) according to the general procedure O. Obtained as a yellow solid (388 mg).

MS (ISP) 510 [(M+H)⁺], 527 [(M+NH₄)⁺] and 532 [(M+Na)⁺]; mp 169 °C.

Example O9

5-tert.-Butoxycarbonylamino-4-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-2-phenylethynyl-benzoic acid methyl ester

- 5 Prepared from 4-amino-5-tert.-butoxycarbonylamino-2-phenylethynyl-benzoic acid methyl ester (Example L14) (396 mg, 1.08 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (273 mg, 1.19 mmol) according to the general procedure O. Obtained as an off-white solid (540 mg).

MS (ISP) 538 [(M+H)⁺], 555 [(M+NH₄)⁺] and 560 [(M+Na)⁺]; mp 158 °C (dec.).

10 Example O10

{2-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-5-morpholin-4-yl-4-phenylethynyl-phenyl}-carbamic acid tert.-butyl ester

- Prepared from (2-amino-5-morpholin-4-yl-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example L12) (483 mg, 1.23 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (422 mg, 1.84 mmol) according to the general procedure O. Obtained as an amorphous orange material (375 mg).

MS (ISP) 565 [(M+H)⁺] and 587 [(M+Na)⁺].

Example O11

- 20 {2-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-5-(2-[2-(2-methoxy-ethoxy)-ethoxy]-ethoxy)-4-phenylethynyl-phenyl}-carbamic acid tert.-butyl ester

Prepared from [2-amino-5-(2-[2-(2-methoxy-ethoxy)-ethoxy]-ethoxy)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example L15) (514 mg, 1.0 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (344 mg, 1.5 mmol) according to the general procedure O. Obtained as a light yellow solid (353 mg).

- 25 MS (ISP) 686 [(M+H)⁺] and 703 [(M+NH₄)⁺]; mp 135-136 °C.

Example O12

{5-tert.-Butoxycarbonylamino-4-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-2-phenylethynyl-phenoxy}-acetic acid tert.-butyl ester

Prepared from (4-amino-5-tert.-butoxycarbonylamino-2-phenylethynyl-phenoxy)-acetic acid tert.-butyl ester (Example L16) (877 mg, 2.0 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (504 mg, 2.2 mmol) according to the general procedure O. Obtained as a yellow solid (723 mg).

5 MS (ISP) 610 [(M+H)⁺], 627 [(M+NH₄)⁺] and 632 [(M+Na)⁺]; mp 95 °C.

Example O13

{5-Cyanomethyl-2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-4-phenylethynyl-phenyl}-carbamic acid tert.-butyl ester

Prepared from (2-amino-5-cyanomethyl-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example L17) (298 mg, 0.86 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (218 mg, 0.95 mmol) according to the general procedure O. Obtained as a yellow solid (299 mg).

MS (ISP) 519 [(M+H)⁺], 536 [(M+NH₄)⁺] and 541 [(M+Na)⁺]; mp 98 °C.

Example O14

15 [2-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-5-(1,4-dioxo-8-aza-spiro[4.5]dec-8-yl)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester

Prepared from [2-amino-5-(1,4-dioxo-8-aza-spiro[4.5]dec-8-yl)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example L18) (393 mg, 0.87 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (301 mg, 1.31 mmol) according to the general procedure O. Obtained as an amorphous orange material (268 mg).

MS (ISP) 621 [(M+H)⁺].

Example O15

[2-[3-(3-Iodo-phenyl)-3-oxo-propionylamino]-5-(2-methoxy-ethoxy)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester

25 Prepared from [2-amino-5-(2-methoxy-ethoxy)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example L11) (840 mg, 2.2 mmol) and 6-(3-iodo-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N7) (780 mg, 2.36 mmol) according to the general procedure O. Obtained as a white solid (1.3 g).

MS (ISP) 655 [(M+H)⁺], 672 [(M+NH₄)⁺], 677 [(M+Na)⁺] and 693 [(M+K)⁺]; mp 172 °C.

Example O16

[2-{3-(3-Cyano-phenyl)-3-oxo-propionylamino}-5-(2,2-dimethyl-[1,3]dioxolan-4-ylmethoxy)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester

Prepared from [2-amino-5-(2,2-dimethyl-[1,3]dioxolan-4-ylmethoxy)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example L20) (439 mg, 1.0 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (345 mg, 1.5 mmol) according to the general procedure O. Obtained as a yellow solid (275 mg).

10 MS (ISP) 610 [(M+H)⁺], 627 [(M+NH₄)⁺] and 632 [(M+Na)⁺]; mp 132-134 °C.

Example O17

5-tert.-Butoxycarbonylamino-4-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-2-iodo-benzoic acid methyl ester

Prepared from 4-amino-5-tert.-butoxycarbonylamino-2-iodo-benzoic acid methyl ester (Example L21) (395 mg, 1.0 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (254 mg, 1.1 mmol) according to the general procedure O. Obtained as an apricot solid (435 mg).

MS (ISP) 564 [(M+H)⁺], 581 [(M+NH₄)⁺] and 586 [(M+Na)⁺]; mp 162-166 °C.

Example O18

20 [[2-{3-(3-Imidazol-1-yl-phenyl)-3-oxo-propionylamino}-5-(2-methoxy-ethoxy)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester.

Prepared from [2-amino-5-(2-methoxy-ethoxy)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example L11) (191 mg, 0.5 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) (135 mg, 0.5 mmol) according to the general procedure O. Obtained as a light brown waxy solid (206 mg).

MS (ISP) 593 [(M-H)⁻]; mp 122-129 °C.

Example O19

[RS]-[2-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-5-(2-oxo-[1,3]dioxolan-4-ylmethoxy)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester

Prepared from [RS]-[2-amino-5-(2-oxo-[1,3]dioxolan-4-ylmethoxy)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example L22) (346 mg, 0.82 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (206 mg, 0.9 mmol) according to the general procedure K. Obtained as a light yellow solid (433 mg).

MS (ISP) 594 [(M-H)⁺]; mp 181 °C.

Example O20

10 {2-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-5-ethoxymethyl-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester

Prepared from (2-amino-5-ethoxymethyl-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example L23) (130 mg, 0.35 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (80 mg, 0.35 mmol) according to the general
15 procedure K. Obtained as an amorphous yellow substance (148 mg).

MS (ISP) 538 [(M-H)⁺].

Example O21

2,2-Dimethyl-propionic acid 5-tert.-butoxycarbonylamino-4-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-2-phenylethynyl-benzyl ester.

20 Prepared from 2,2-dimethyl-propionic acid 4-amino-5-tert.-butoxycarbonylamino-2-phenylethynyl-benzyl ester (Example L24) (155 mg, 0.37 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (94 mg, 0.41 mmol) according to the general procedure K. Obtained as an amorphous light orange substance (184 mg).

MS (ISP) 611 [(M+NH₄)⁺].

25 Example O22

(RS)-[2-[3-(3-Imidazol-1-yl-phenyl)-3-oxo-propionylamino]-4-phenylethynyl-5-(tetrahydro-pyran-2-yloxymethyl)-phenyl]-carbamic acid tert.-butyl ester.

Prepared from (RS)-[2-amino-4-phenylethynyl-5-(tetrahydro-pyran-2-yloxymethyl)-phenyl]-carbamic acid tert.-butyl ester (Example L25) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) according to the general procedure O. Obtained as an amorphous yellow substance (267 mg).

5 MS (ISP) 635 [(M+H)⁺].

Example O23

[2-[3-(3-Imidazol-1-yl-phenyl)-3-oxo-propionylamino]-5-(4-methoxy-piperidin-1-yl)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester.

Prepared from [2-amino-5-(4-methoxy-piperidin-1-yl)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example L26) (236 mg, 0.56 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) (151 mg, 0.56 mmol) according to the general procedure O. Obtained as an amorphous yellow substance (253 mg).

MS (ISP) 634 [(M+H)⁺].

15 Example O24

[2-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-5-(4-methoxy-piperidin-1-yl)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester.

Prepared from [2-amino-5-(4-methoxy-piperidin-1-yl)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example L26) (224 mg, 0.53 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (125 mg, 0.53 mmol) according to the general procedure O. Obtained as a yellow foam (274 mg).

MS (ISN) 591 [(M-H)⁻]; mp 97-100 °C.

Example O25

[5-Cyanomethoxy-2-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester.

Prepared from (2-amino-5-cyanomethoxy-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example L27) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) according to the general procedure O. Obtained as an amorphous yellow substance (169 mg).

MS (ISP) 576 [(M+H)⁺].

Example O26

(5-Cyanomethoxy-2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester.

- 5 Prepared from (2-amino-5-cyanomethoxy-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example L27) (80 mg, 0.22 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (55 mg, 0.24 mmol) according to the general procedure O. Obtained as an amorphous yellow substance (96 mg).

MS (ISN) 533 [(M-H)⁻].

10 Example O27

(RS)-[4-(4-Fluoro-phenylethynyl)-2-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-5-(tetrahydro-pyran-2-yloxymethyl)-phenyl]-carbamic acid tert.-butyl ester.

- Prepared from (RS)-[2-amino-4-(4-fluoro-phenylethynyl)-5-(tetrahydro-pyran-2-yloxymethyl)-phenyl]-carbamic acid tert.-butyl ester (Example L28) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) according to the general procedure O. Obtained as an amorphous yellow substance (990 mg).

MS (ISN) 651 [(M-H)⁻].

Example O28

- 20 (RS)-[4-(4-Fluoro-phenylethynyl)-2-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-5-{methyl-[2-(tetrahydro-pyran-2-yloxy)-ethyl]-amino}-phenyl]-carbamic acid tert.-butyl ester

- Prepared from (RS)-[2-amino-4-(4-fluoro-phenylethynyl)-5-{methyl-[2-(tetrahydro-pyran-2-yloxy)-ethyl]-amino}-phenyl]-carbamic acid tert.-butyl ester (Example L29) (242 mg, 0.5 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) (135 mg, 0.5 mmol) according to the general procedure O. Obtained as an amorphous yellow substance (198 mg).

MS (ISN) 694 [(M-H)⁻].

Example O29

{5-(4,4-Diethoxy-piperidin-1-yl)-2-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-4-phenylethynyl-phenyl}-carbamic acid tert.-butyl ester

- Prepared from [2-amino-5-(4,4-diethoxy-piperidin-1-yl)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example L18) (321 mg, 0.67 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) (240 mg, 0.89 mmol) according to the general procedure O. Obtained as an orange foam (295 mg).

MS (ISP) 692 [(M+H)⁺].

Example O30

- 10 (RS)-{2-[3-(3-Imidazol-1-yl-phenyl)-3-oxo-propionylamino]-4-phenylethynyl-5-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-phenyl}-carbamic acid tert.-butyl ester

- Prepared from (RS)-{2-amino-4-phenylethynyl-5-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-phenyl}-carbamic acid tert.-butyl ester (Example L30) (346 mg, 0.76 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) (300 mg, 1.11 mmol) according to the general procedure O. Obtained as a yellow foam (196 mg).

MS (ISP) 665 [(M+H)⁺].

Example O31

(RS)-{4'-Fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-2-[4-(tetrahydro-pyran-2-yloxy)-piperidin-1-yl]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- 20 Prepared from (RS)-[2-amino-4-(4-fluoro-phenylethynyl)-5-(tetrahydro-pyran-2-yloxymethyl)-phenyl]-carbamic acid tert.-butyl ester (Example L28) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) according to the general procedure O. Obtained as a yellow solid (282 mg).

MS (ISP) 698 [(M+H)⁺]; mp 129-132 °C.

- 25 Example O32

{5-(2-tert.-Butoxy-ethoxy)-4-(4-fluoro-phenylethynyl)-2-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-phenyl}-carbamic acid tert.-butyl ester

Prepared from [2-amino-5-(2-tert.-butoxy-ethoxy)-4-(4-fluoro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester (Example L32) (560 mg, 1.27 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) (413 mg, 1.53 mmol) according to the general procedure O. Obtained as a yellow solid (507 mg).

- 5 MS (ISP) 655 [(M+H)⁺]; mp 62-65 °C.

Example O33

(RS)-(4'-Fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-2-{4-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-piperidin-1-yl}-biphenyl-4-yl)-carbamic acid tert.-butyl ester

- 10 Prepared from (RS)-(5-amino-4'-fluoro-2-{4-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-piperidin-1-yl}-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L33) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) according to the general procedure O. Obtained as a yellow solid (473 mg).

MS (ISP) 742 [(M+H)⁺]; mp 57-58 °C.

- 15 Example O34

(RS)-[4'-Fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-2-(tetrahydro-pyran-2-yloxymethyl)-biphenyl-4-yl]-carbamic acid tert.-butyl ester

- Prepared from (RS)-[5-amino-4'-fluoro-2-(tetrahydro-pyran-2-yloxymethyl)-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example L34) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) according to the general procedure O.
20 Obtained as a light yellow solid (330 mg).

Example O35

{2-Cyanomethoxy-4'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- 25 Prepared from (5-amino-2-cyanomethoxy-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L35) (190 mg, 0.53 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) (164 mg, 0.61 mmol) according to the general procedure O. Obtained as a yellow gum (90 mg).

MS (ISP) 570 [(M+H)⁺].

Example O36

{2-Dimethylaminomethyl-4'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (5-amino-2-dimethylaminomethyl-4'-fluoro-biphenyl-4-yl)-carbamic acid
5 tert.-butyl ester (Example L36) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) according to the general procedure O. Obtained as a light yellow solid (329 mg).

Example O37

{2-(2,2-Dimethyl-tetrahydro-[1,3]dioxolo[4,5-c]pyrrol-5-yl)-4'-fluoro-5-[3-(3-imidazol-
10 1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from [5-amino-2-(2,2-dimethyl-tetrahydro-[1,3]dioxolo[4,5-c]pyrrol-5-yl)-4'-fluoro-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example L37) (465 mg, 1.05 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) (314 mg, 1.16 mmol) according to the general procedure O. Obtained as an amorphous yellow
15 substance (540 mg).

MS (ISN) 654 [(M-H)⁺].

Example O38

{4'-Fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-2-methoxy-biphenyl-4-yl}-carbamic acid tert.-butyl ester

20 Prepared from (5-amino-4'-fluoro-2-methoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L38) (332 mg, 1.0 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) (270 mg, 1.0 mmol) according to the general procedure O. Obtained as an amorphous brown substance (328 mg).

MS (ISN) 543 [(M-H)⁺].

25 Example O39

{2-(1,4-Dioxa-8-aza-spiro[4.5]dec-8-yl)-4'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from [5-amino-2-(1,4-dioxo-8-aza-spiro[4.5]dec-8-yl)-4'-fluoro-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example L39) (444 mg, 1.0 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) (321 mg, 1.19 mmol) according to the general procedure O. Obtained as a brown solid (457 mg).

- 5 MS (ISN) 654 [(M-H)⁺]; mp 110-115 °C (dec.).

Example O40

{4'-Fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-2-methyl-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- Prepared from (5-amino-4'-fluoro-2-methyl-biphenyl-4-yl)-carbamic acid tert.-butyl ester
10 (Example L40) (316 mg, 1.0 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) (297 mg, 1.1 mmol) according to the general procedure O. Obtained as an amorphous yellow substance (361 mg).

MS (ISP) 529 [(M+H)⁺].

Example O41

- 15 {4-tert.-Butoxycarbonylamino-4'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-2-yloxy}-acetic acid tert.-butyl ester

- Prepared from (5-amino-4-tert.-butoxycarbonylamino-4'-fluoro-biphenyl-2-yloxy)-acetic acid tert.-butyl ester (Example L41) (1.4 g, 3.24 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) (885 mg, 3.27 mmol) according to the
20 general procedure O. Obtained as a light brown solid (759 mg).

MS (ISP) 645 [(M+H)⁺]; mp 82-85 °C.

Example O42

{2-Chloro-4'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- 25 Prepared from (5-amino-2-chloro-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L42) (168 mg, 0.5 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) (252 mg, 0.93 mmol) according to the general procedure O. Obtained as a light yellow solid (156 mg).

MS (ISN) 547 [(M-H)⁺].

Example O43

[4'-Fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-2-(2-methoxy-ethoxy)-biphenyl-4-yl]-carbamic acid tert.-butyl ester

- 5 Prepared from [5-amino-4'-fluoro-2-(2-methoxy-ethoxy)-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example L43) (188 mg, 0.5 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) (260 mg, 0.96 mmol) according to the general procedure O. Obtained as an orange solid (218 mg).

MS (ISP) 589 [(M+H)⁺]; mp 61-63 °C.

10 Example O44

{2-(2-tert.-Butoxy-ethoxy)-4'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- Prepared from [5-amino-2-(2-tert.-butoxy-ethoxy)-4'-fluoro-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example L44) (209 mg, 0.5 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) (135 mg, 0.5 mmol) according to the
15 general procedure O. Obtained as a beige solid (194 mg).

MS (ISP) 631 [(M+H)⁺]; mp 101 °C.

Example O45

- [4'-Fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-2-(2-oxo-oxazolidin-3-yl)-biphenyl-4-yl]-carbamic acid tert.-butyl ester
20

Prepared from [5-amino-4'-fluoro-2-(2-oxo-oxazolidin-3-yl)-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example L45) (130 mg, 0.34 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) (92 mg, 0.34 mmol) according to the general procedure O. Obtained as an amorphous yellow substance (86 mg).

- 25 MS (ISP) 600 [(M+H)⁺].

Example O46

{4'-Fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-2-methoxy-2'-methyl-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (5-amino-4'-fluoro-2-methoxy-2'-methyl-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L46) (173 mg, 0.5 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) (199 mg, 1.47 mmol) according to the general procedure O. Obtained as an orange solid (182 mg).

- 5 MS (ISP) 559 [(M+H)⁺]; mp 99-102 °C.

Example O47

{2-tert.-Butoxy-4'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- 10 Prepared from (5-amino-2-tert.-butoxy-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L47) (187 mg, 0.5 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) (135 mg, 0.5 mmol) according to the general procedure O. Obtained as an amorphous yellow substance (237 mg).

MS (ISN) 585 [(M-H)⁻].

Example O48

- 15 {2-tert.-Butoxy-2'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- Prepared from (5-amino-2-tert.-butoxy-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L48) (187 mg, 0.5 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) (135 mg, 0.5 mmol) according to the general procedure
20 O. Obtained as an amorphous yellow substance (234 mg).

MS (ISN) 585 [(M-H)⁻].

Example O49

(RS)-[4'-Fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-2-[(R)-3-(tetrahydro-pyran-2-yloxy)-pyrrolidin-1-yl]-biphenyl-4-yl]-carbamic acid tert.-butyl ester

- 25 Prepared from (RS)-[5-amino-4'-fluoro-2-[(R)-3-(tetrahydro-pyran-2-yloxy)-pyrrolidin-1-yl]-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example L49) (236 mg, 0.5 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) (200 mg, 0.74 mmol) according to the general procedure O. Obtained as an orange solid (188 mg).

MS (ISP) 684 [(M+H)⁺]; mp 99-103 °C.

Example O50

{2'-Fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-2-methoxy-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- 5 Prepared from (5-amino-2'-fluoro-2-methoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L50) (159 mg, 0.5 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) (135 mg, 0.5 mmol) according to the general procedure O. Obtained as an amorphous yellow substance (199 mg).

MS (ISP) 543 [(M-H)⁻].

10 Example O51

{2-tert.-Butoxy-5-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-4'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- Prepared from (5-amino-2-tert.-butoxy-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L47) (187 mg, 0.5 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (126 mg, 0.55 mmol) according to the general procedure O.
15 Obtained as an amorphous light pink substance (196 mg).

MS (ISP) 546 [(M+H)⁺].

Example O52

- 20 {2-tert.-Butoxy-5-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-2'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (5-amino-2-tert.-butoxy-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L48) (187 mg, 0.5 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (126 mg, 0.55 mmol) according to the general procedure O. Obtained as an amorphous light pink substance (197 mg).

- 25 MS (ISP) 546 [(M+H)⁺].

Example O53

{5-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-2'-fluoro-2-methoxy-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (5-amino-2'-fluoro-2-methoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L50) (112 mg, 0.35 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (80 mg, 0.35 mmol) according to the general procedure O. Obtained as a yellow foam (131 mg).

- 5 MS (ISP) 502 [(M-H)⁺].

Example O54

(2'-Fluoro-2-methoxy-5-{3-[3-(2-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino}-biphenyl-4-yl)-carbamic acid tert.-butyl ester

- 10 Prepared from (5-amino-2'-fluoro-2-methoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L50) (249 mg, 0.75 mmol) and 3-[3-(2-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester (Example M8) (275 mg, 0.92 mmol) according to the general procedure O. Obtained as a yellow solid (312 mg).

MS (ISP) 559 [(M+H)⁺]; mp 83-86 °C.

Example O55

- 15 {5-[3-(5-Cyano-thiophen-2-yl)-3-oxo-propionylamino]-2'-fluoro-2-methoxy-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- Prepared from (5-amino-2'-fluoro-2-methoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L50) (166 mg, 0.5 mmol) 3-(5-cyano-thiophen-2-yl)-3-oxo-propionic acid tert.-butyl ester (Example M17) (138 mg, 0.55 mmol) according to the general procedure
20 O. Obtained as a light yellow solid (244 mg).

MS (ISP) 510 [(M+H)⁺]; mp 200 °C (dec.).

Example O56

{2'-Fluoro-2-methoxy-5-[3-oxo-3-(3-[1,2,4]triazol-4-yl-phenyl)-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- 25 Prepared from (5-amino-2'-fluoro-2-methoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L50) (166 mg, 0.5 mmol) and 3-oxo-3-(3-[1,2,4]triazol-4-yl-phenyl)-propionic acid ethyl ester (Example M1) (260 mg, 1.0 mmol) according to the general procedure O. Obtained as a yellow gum (70 mg).

MS (ISP) 546 [(M+H)⁺].

Example O57

{5-[3-(2-Cyano-pyridin-4-yl)-3-oxo-propionylamino]-2'-fluoro-2-methoxy-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- 5 Prepared from (5-amino-2'-fluoro-2-methoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L50) and 3-(2-cyano-pyridin-4-yl)-3-oxo-propionic acid tert.-butyl ester (Example M10) according to the general procedure O. Obtained as a light yellow solid (189 mg).

MS (ISP) 522 [(M+NH₄)⁺].

10 Example O58

(2-tert.-Butoxy-4'-fluoro-5-[3-[3-(2-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino]-biphenyl-4-yl)-carbamic acid tert.-butyl ester

- Prepared from (5-amino-2-tert.-butoxy-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L47) (140 mg, 0.37 mmol) and 3-[3-(2-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester (Example M8) (111 mg, 0.37 mmol) according to the
15 general procedure O. Obtained as an amorphous yellow substance (139 mg).

MS (ISP) 601 [(M+H)⁺].

Example O59

- {5-[3-(5-Cyano-2-fluoro-phenyl)-3-oxo-propionylamino]-2'-fluoro-2-methoxy-biphenyl-4-yl}-carbamic acid tert.-butyl ester
20

Prepared from (5-amino-2'-fluoro-2-methoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L50) (166 mg, 0.5 mmol) and 3-(5-cyano-2-fluoro-phenyl)-3-oxo-propionic acid ethyl ester (Example M18) (141 mg, 0.6 mmol) according to the general procedure O. Obtained as an amorphous yellow substance (165 mg).

- 25 MS (ISP) 522 [(M+H)⁺].

Example O60

{2-tert.-Butoxy-5-[3-(2-cyano-pyridin-4-yl)-3-oxo-propionylamino]-4'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (5-amino-2-tert.-butoxy-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L47) (140 mg, 0.37 mmol) and 3-(2-cyano-pyridin-4-yl)-3-oxo-propionic acid tert.-butyl ester (Example M10) (91 mg, 0.37 mmol) according to the general procedure O. Obtained as an amorphous yellow substance (164 mg).

- 5 MS (ISP) 547 [(M+H)⁺].

Example O61

{2-tert.-Butoxy-2'-fluoro-5-[3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- 10 Prepared from (5-amino-2-tert.-butoxy-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L48) (187 mg, 0.5 mmol) and 3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionic acid ethyl ester (Example M2) (180 mg, 0.69 mmol) according to the general procedure O. Obtained as a light yellow solid (257 mg).

MS (ISP) 588 [(M+H)⁺]; mp 47-50 °C.

Example O62

- 15 [5-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-2-(1,4-dioxo-8-aza-spiro[4.5]dec-8-yl)-2'-fluoro-biphenyl-4-yl]-carbamic acid tert.-butyl ester

- Prepared from [5-amino-2-(1,4-dioxo-8-aza-spiro[4.5]dec-8-yl)-2'-fluoro-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example L51) (222 mg, 0.5 mmol) and 3-(3-cyano-phenyl)-3-oxo-propionic acid tert.-butyl ester (Example M3) (182 mg, 0.8 mmol)
20 according to the general procedure O. Obtained as an amorphous yellow substance (258 mg).

MS (ISP) 615 [(M+H)⁺].

Example O63

- 25 {2-(1,4-Dioxo-8-aza-spiro[4.5]dec-8-yl)-2'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from [5-amino-2-(1,4-dioxo-8-aza-spiro[4.5]dec-8-yl)-2'-fluoro-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example L51) (222 mg, 0.5 mmol) and 6-(3-imidazol-1-yl-phenyl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N10) (135 mg, 0.5 mmol) according to the general procedure O. Obtained as an amorphous yellow substance (294 mg).

MS (ISP) 656 [(M+H)⁺].

Example O64

(2-tert.-Butoxy-2'-fluoro-5-[3-[3-(2-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino]-biphenyl-4-yl)-carbamic acid tert.-butyl ester

- 5 Prepared from (5-amino-2-tert.-butoxy-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L48) (187 mg, 0.5 mmol) and 2,2-dimethyl-6-[3-(2-methyl-imidazol-1-yl)-phenyl]-[1,3]dioxin-4-one (Example N15) (142 mg, 0.5 mmol) according to the general procedure O. Obtained as an amorphous yellow substance (234 mg).

MS (ISP) 601 [(M+H)⁺].

10 Example O65

{2-tert.-Butoxy-5-[3-(2-cyano-pyridin-4-yl)-3-oxo-propionylamino]-2'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- Prepared from (5-amino-2-tert.-butoxy-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L48) (187 mg, 0.5 mmol) and 4-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-pyridine-2-carbonitrile (Example N16) (115 mg, 0.5 mmol) according to the general procedure O. Obtained as an amorphous yellow substance (216 mg).
- 15

MS (ISP) 547 [(M+H)⁺].

Example O66

- (2-tert.-Butoxy-2'-fluoro-5-[3-[3-(2-methylsulfanyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino]-biphenyl-4-yl)-carbamic acid tert.-butyl ester
- 20

Prepared from (5-amino-2-tert.-butoxy-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L48) (187 mg, 0.5 mmol) and 3-[3-(2-methylsulfanyl-imidazol-1-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester (Example M13) (211 mg, 0.63 mmol) according to the general procedure O. Obtained as a light yellow solid (260 mg).

- 25 MS (ISN) 631 [(M-H)⁻]; mp 59-62 °C.

Example O67

{5-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-2',5'-difluoro-2-methoxy-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (5-amino-2',5'-difluoro-2-methoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L52) (175 mg, 0.5 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (115 mg, 0.5 mmol) according to the general procedure O. Obtained as an amorphous yellow substance (136 mg).

5 MS (ISP) 522 [(M+H)⁺].

Example O68

{2',5'-Difluoro-2-methoxy-5-[3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (5-amino-2',5'-difluoro-2-methoxy-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L52) (175 mg, 0.5 mmol) and 3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionic acid ethyl ester (Example M2) (130 mg, 0.5 mmol) according to the general procedure O. Obtained as an amorphous light yellow substance (185 mg).

MS (ISN) 562 [(M-H)⁻].

Example O69

15 {2-tert.-Butoxy-5-[3-(3-cyano-thiophen-2-yl)-3-oxo-propionylamino]-2'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester

Prepared from (5-amino-2-tert.-butoxy-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L48) (187 mg, 0.5 mmol) and 6-(3-cyano-thiophen-2-yl)-2,2-dimethyl-[1,3]dioxin-4-one (Example N3) (130 mg, 0.55 mmol) according to the general procedure
20 O. Obtained as a yellow oil (278 mg).

MS (ISN) 550 [(M-H)⁻].

Example O70

{2-tert.-Butoxy-5-[3-(5-cyano-thiophen-2-yl)-3-oxo-propionylamino]-2'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester

25 Prepared from (5-amino-2-tert.-butoxy-2'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L48) (187 mg, 0.5 mmol) 3-(5-cyano-thiophen-2-yl)-3-oxo-propionic acid tert.-butyl ester (Example M17) (138 mg, 0.55 mmol) according to the general procedure O. Obtained as an amorphous yellow substance (268 mg).

MS (ISN) 550 [(M-H)⁻].

Example O71

{2-tert.-Butoxy-4'-fluoro-5-[3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester

- 5 Prepared from (5-amino-2-tert.-butoxy-4'-fluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L47) (187 mg, 0.5 mmol) and 3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionic acid ethyl ester (Example M2) (156 mg, 0.6 mmol) according to the general procedure O. Obtained as a yellow gum (198 mg).

MS (ISP) 588 [(M+H)⁺].

10 Example O72

[5-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-2'-fluoro-2-(2-methoxy-ethoxy)-biphenyl-4-yl]-carbamic acid tert.-butyl ester

- Prepared from [5-amino-2'-fluoro-2-(2-methoxy-ethoxy)-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example L53) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) according to the general procedure O. Obtained as a yellow solid (188 mg).

MS (ISP) 548 [(M+H)⁺].

Example O73

- 20 (RS)-[5-[3-(3-Cyano-phenyl)-3-oxo-propionylamino]-2'-fluoro-2-(tetrahydro-pyran-2-yloxymethyl)-biphenyl-4-yl]-carbamic acid tert.-butyl ester

Prepared from (RS)-[5-amino-2'-fluoro-2-(tetrahydro-pyran-2-yloxymethyl)-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example L54) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) according to the general procedure O. Obtained as a yellow solid (155 mg).

- 25 MS (ISP) 548 [(M+NH₄)⁺].

Example O74

(2'-Fluoro-2-(4-methoxy-benzyloxy)-5-[3-[3-(3-methyl-isoxazol-5-yl)-phenyl]-3-oxo-propionylamino]-biphenyl-4-yl)-carbamic acid tert.-butyl ester

Prepared [5-amino-2'-fluoro-2-(4-methoxy-benzyloxy)-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example L55) (438 mg, 1.0 mmol) and 3-[3-(3-methyl-isoxazol-5-yl)-phenyl]-3-oxo-propionic acid tert.-butyl ester (Example M14) (301 mg, 1.0 mmol) according to the general procedure O. Obtained as an amorphous light yellow substance
5 (561 mg).

MS (ISP) 666 [(M+H)⁺].

Example O75

{2-tert.-Butoxy-5-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-2',5'-difluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester

10 Prepared from (5-amino-2-tert.-butoxy-2',5'-difluoro-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L56) (196 mg, 0.5 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (115 mg, 0.5 mmol) according to the general procedure O. Obtained as an amorphous beige substance (155 mg).

MS (ISP) 564 [(M+H)⁺].

15 Example O76

{5-tert.-Butoxy-4-(4-fluoro-phenylethynyl)-2-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-phenyl}-carbamic acid tert.-butyl ester

Prepared from [2-amino-5-tert.-butoxy-4-(4-fluoro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester (Example L57) (160 mg, 0.4 mmol) and 3-(3-imidazol-1-yl-phenyl)-
20 3-oxo-propionic acid tert.-butyl ester (Example M4) (115 mg, 0.4 mmol) according to the general procedure O. Obtained as an amorphous yellow substance (140 mg).

MS (ISP) 611 [(M+H)⁺].

Example O77

{5-tert.-Butoxy-4-(4-fluoro-phenylethynyl)-2-[3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionylamino]-phenyl}-carbamic acid tert.-butyl ester
25

Prepared from [2-amino-5-tert.-butoxy-4-(4-fluoro-phenylethynyl)-phenyl]-carbamic acid tert.-butyl ester (Example L57) (160 mg, 0.4 mmol) and 3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionic acid ethyl ester (Example M2) (104 mg, 0.4 mmol) according to the general procedure O. Obtained as a yellow gum (150 mg).

MS (ISP) 612 [(M+H)⁺].

General procedure P:

Preparation of 4,8-diaryl-1,3-dihydro-benzo[b][1,4]diazepin-2-ones, 4-aryl-8-aryloxy-1,3-dihydro-benzo[b][1,4]diazepin-2-ones or 4-aryl-8-arylethynyl-1,3-dihydro-

5 benzo[b][1,4]diazepin-2-ones

A suspension of the {2-[3-aryl-3-oxo-propionylamino]-4-aryl-phenyl}-carbamic acid tert-butyl ester or {2-[3-aryl-3-oxo-propionylamino]-4-arylethynyl-phenyl}-carbamic acid tert-butyl ester (1.0 mmol) in CH₂Cl₂ (5 mL) [anisole or 1,3-dimethoxybenzene (5-15 mmol) can be added if necessary] was treated with TFA (0.5-5.0 mL) at 0 °C and stirring was
10 continued at 23 °C until tlc indicated complete consumption of the starting material. The solvent was removed in vacuum, the residue treated with little ether, whereupon it crystallized. The solid was stirred with sat. NaHCO₃-sol., filtered, washed with H₂O and ether or mixtures of ether/hexane and was dried to give the title compound, which if necessary can be purified by crystallization from THF/CH₂Cl₂/ether/hexane.

15 General procedure Q:

Preparation of 4-[3-(Amino-4-carbonyl)-phenyl]-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-ones by Pd-catalyzed carbonylative amination of 4-(3-iodophenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

A solution of the 4-(3-iodophenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-
20 2-one (1.0 mmol), the secondary amine (5.0 mmol), PPh₃ (6 mol%) or dppp (3 mol%), Pd(OAc)₂ (3 mol%) and Et₃N (2.0 mmol) in DMF (4 mL) was stirred at 23 °C under CO-atmosphere until tlc indicated complete consumption of the iodide. After dilution with EtOAc, washing with sat. NaHCO₃-sol. and brine, the organic phase was dried over Na₂SO₄. Removal of the solvent left a brown oil, which was purified by silica gel column
25 chromatography with hexane/EtOAc to give the title compound.

General procedure R:

Preparation of (5-hydroxy-2-nitro-phenyl)-carbamic acid tert.-butyl esters by Rh-catalyzed deallylation of (5-allyloxy-2-nitro-phenyl)-carbamic acid tert.-butyl esters

A mixture of the (5-allyloxy-2-nitro-phenyl)-carbamic acid tert.-butyl ester, (PPh₃)₃RhCl
30 (5 mol%) and DABCO (20 mol%) in EtOH was refluxed for 2.5 h according to *J. Org. Chem.* 1973, 38, 3224. Added 5% citric acid, stirred at 23 °C for 15 min, extracted with EtOAc, washed with brine, dried over MgSO₄. Removal of the solvent left an orange solid, which was purified by silica gel column chromatography with hexane/EtOAc to give the title compound.

Example R1

(5-Hydroxy-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester

Prepared from (5-allyloxy-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example B4), $(\text{PPh}_3)_3\text{RhCl}$ (5 mol%) and DABCO (20 mol%) in EtOH according to the general procedure R. Obtained as a yellow solid.

MS (ISN) 379 $[(\text{M}-\text{H})^-]$; mp 140 °C.

General procedure S:

- Preparation of 5-O-substituted-(4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl esters from (5-hydroxy-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester .
- A mixture of the (5-hydroxy-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example R1), KHCO_3 and the appropriate alkylating reagent were stirred in DMF at 23 to 60 °C until tlc indicated complete conversion. Dilution with EtOAc was followed by aqueous workup with 5% citric acid, sat. NaHCO_3 -sol., brine and drying over MgSO_4 .
- Removal of the solvent left a crude material, which was purified by silica gel column chromatography with hexane/EtOAc to give the title compound.

Example S1

(5-tert.-Butoxycarbonylamino-2-iodo-4-nitro-phenoxy)-acetic acid tert.-butyl ester

- Prepared from (5-hydroxy-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example R1) (1.23 g, 3.24 mmol), KHCO_3 (0.39 g, 3.89 mmol) and tert.-butyl bromoacetate (0.59 mL, 3.89 mmol) according to the general procedure S. Obtained as a yellow solid (1.5 g, 94 %).

MS (ISP) 495 $[(\text{M}+\text{H})^+]$, 512 $[(\text{M}+\text{NH}_4)^+]$ and 517 $[(\text{M}+\text{Na})^+]$; mp 103 °C.

Example S2

- (5-Cyanomethoxy-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester

Prepared from (5-hydroxy-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example R1) (614 mg, 1.62 mmol), KHCO_3 (208 mg, 2.08 mmol) and bromoacetonitrile (0.21 mL, 3.16 mmol) according to the general procedure S. Obtained as a yellow solid (574 mg, 85 %).

MS (ISN) 418 [(M-H)⁻]; mp 125 °C.

Example S3

(RS)-{4-Iodo-2-nitro-5-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-phenyl}-carbamic acid tert.-butyl ester

- 5 Prepared from (5-hydroxy-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example R1) (760 mg, 2 mmol), KHCO₃ (260 mg, 4 mmol) and 2-(2-bromoethoxy)tetrahydro-2H-pyran (0.6 mL, 2.6 mmol) according to the general procedure S. Obtained as an orange oil (804 mg, 79 %).

MS (EI) 508 (M⁺).

10 Example S4

(5-tert.-Butoxy-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester

- N,N-Dimethylformamide di-tert.butylacetal (19.2 mL, 80 mmol) was added dropwise within 15 min to a solution of (5-hydroxy-4-iodo-2-nitro-phenyl)-carbamic acid tert.-butyl ester (Example R1) (7.60 g, 20 mmol) in toluene at 80 °C and stirring was continued
15 at 80 °C for 3 h (cf. *Synthesis* 1983, 135). Obtained as a yellow solid (5.97 g, 68 %).

MS (ISN) 435 [(M-H)⁻]; mp 94 °C.

Example 1

3-(7-Iodo-4-oxo-8-thiomorpholin-4-yl)-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

- 20 Prepared from 2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-4-iodo-5-thiomorpholin-4-yl-phenyl}-carbamic acid tert.-butyl ester (Example O1) (629 mg, 1.04 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as an olive solid (437 mg).

mp 227-228 °C (dec.).

25 Example 2

3-(7-Iodo-8-morpholin-4-yl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

Prepared from {2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-4-iodo-5-morpholin-4-yl-phenyl}-carbamic acid tert.-butyl ester (Example O2) (518 mg, 0.877 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a beige solid (309 mg).

- 5 MS (EI) 472 (M⁺); mp 224 °C (dec.).

Example 3

3-(8-Chloro-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

- 10 Prepared from (2-amino-5-chloro-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example L3) (171 mg, 0.5 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (183 mg, 0.6 mmol) according to the general procedure O. Obtained as a light yellow solid (284 mg). This material was deprotected and cyclized by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as an orange solid (483 mg).

- 15 MS (ISP) 343 [(M+H)⁺] and 345 [(M+2+Na)⁺]; mp 248-251 °C (dec.).

Example 4

3-(8-Methyl-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

- 20 Prepared from (2-amino-5-methyl-4-phenylethynyl-phenyl)-carbamic acid tert.-butyl ester (Example L4) (161 mg, 0.5 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (230 mg, 0.75 mmol, 75% pure) according to the general procedure O. Obtained as a light yellow solid (227 mg). This material was deprotected and cyclized by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a light yellow solid (83 mg).

- 25 MS (ISP) 375 (M⁺); mp 237-239 °C (dec.).

Example 5

3-[8-(4-Methyl-piperazin-1-yl)-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from [2-amino-5-(4-methyl-piperazin-1-yl)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example L5) (203 mg, 0.5 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (230 mg, 0.75 mmol, 75% pure) according to the general procedure O. Obtained by chromatography as an orange oil (181 mg). This material was deprotected and cyclized by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as an orange solid (82 mg).

MS (ISP) 460.5 [(M+H)⁺]; mp 222-224 °C (dec.).

Example 6

3-[8-(1,1-Dioxo-thiomorpholin-4-yl)-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from [2-amino-5-(1,1-dioxo-6-thiomorpholin-4-yl)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example L19) (220 mg, 0.5 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (172 mg, 0.75 mmol) according to the general procedure O. The obtained material was deprotected and cyclized by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (47 mg).

MS (ISP) 495 [(M+H)⁺]; mp >250 °C (dec.).

Example 7

3-(8-Chloro-4-oxo-7-phenyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

Prepared from {2-chloro-5-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O3) (720 mg, 1.47 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as an off-white solid (457 mg).

MS (EI) 371 (M⁺) and 373 [(M+2)⁺]; mp 242-244 °C (dec.).

Example 8

3-(8-Methyl-4-oxo-7-phenyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

Prepared from (5-amino-2-methyl-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example L7) (298 mg, 1.0 mmol) and 3-(2,2-dimethyl-6-oxo-6H-[1,3]dioxin-4-yl)-benzonitrile (Example N4) (460 mg, 1.5 mmol) according to the general procedure O. The obtained

material (351 mg) was deprotected and cyclized by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a light yellow solid (206 mg).

MS (EI) 351 (M⁺); mp 236-239 °C (dec.).

Example 9

- 5 3-[8-(1,4-Dioxa-8-aza-spiro[4.5]dec-8-yl)-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from [2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-5-(1,4-dioxa-8-aza-spiro[4.5]dec-8-yl)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example O14) (265 mg, 0.43 mmol) by treatment with TFA in CH₂Cl₂ according to the general
10 procedure P. Obtained as a brown solid (121 mg).

MS (ISP) 503 [(M+H)⁺]; mp 239-243 °C (dec.).

Example 10

3-(8-Morpholin-4-yl-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

- 15 Prepared from [2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-5-morpholin-4-yl-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester O10 (370 mg, 0.66 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a light brown solid (216 mg).

MS (EI) 446 (M⁺); mp 239-243 °C (dec.).

20 Example 11

3-[8-(2-Dimethylamino-ethylsulfanyl)-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

- Prepared from [2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-5-(2-dimethylamino-ethylsulfanyl)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (example O4) (166
25 mg, 0.28 mmol) by treatment with TFA and anisole in CH₂Cl₂ according to the general procedure P. Obtained as a light yellow solid (103 mg).

MS (ISP) 465 [(M+H)⁺]; mp 197 °C (dec.).

Example 12

[4-(3-Cyano-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-ylsulfanyl]-acetic acid methyl ester

- Prepared from {5-tert.-butoxycarbonylamino-4-[3-(3-cyano-phenyl)-3-oxo-
5 propionylamino]-2-phenylethynyl-phenylsulfanyl]-acetic acid methyl ester (Example
O5)(421 mg, 0.72 mmol) by treatment with TFA and anisole in CH₂Cl₂ according to the
general procedure P. Obtained as a light yellow solid (309 mg).

MS (ISP) 465 [(M+H)⁺]; mp 201 °C (dec.).

Example 13

- 10 [4-(3-Cyano-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-ylsulfanyl]-acetic acid

- A solution of [4-(3-cyano-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-ylsulfanyl]-acetic acid methyl ester (Example 12) (265 mg, 0.57
mmol) and LiOH·H₂O (26 mg, 0.63 mmol) in THF (5 mL), MeOH (1 mL) and H₂O (1
15 mL) was stirred at 23 °C for 24 h. Obtained as a light yellow solid (257 mg).

MS (ISP) 452 [(M+H)⁺]; mp 202 °C (dec.).

Example 14

[4-(3-Cyano-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-yl]-acetic acid methyl ester

- 20 Prepared from {5-tert.-butoxycarbonylamino-4-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-2-phenylethynyl-phenyl}-acetic acid methyl ester (Example O6) (846 mg, 1.53 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a light yellow solid (557 mg).

MS (EI) 433 (M⁺); mp 236 °C (dec.).

- 25 Example 15

[4-(3-Cyano-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-yl]-acetic acid

A solution of LiOH·H₂O (54 mg, 1.28 mmol) in H₂O (2 mL) and MeOH (2 mL) was added to [4-(3-cyano-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-yl]-acetic acid methyl ester (Example 14) (505 mg, 1.17 mmol) in THF (10 mL) and the reaction mixture was stirred at 23 °C for 48 h. Obtained as a light yellow solid (62 mg).

- 5 MS (ISP) 452 [(M+H)⁺]; mp 248 °C (dec.).

Example 16

4-(3-Cyano-phenyl)-8-iodo-2-oxo-2,3-dihydro-1H-benzo[b][1,4]diazepine-7-carboxylic acid methyl ester

- Prepared from 5-tert.-butoxycarbonylamino-4-[3-(3-cyano-phenyl)-3-oxo-
10 propionylamino]-2-iodo-benzoic acid methyl ester (Example O17) (430 mg, 0.763mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a salmon colored solid (199 mg).

MS (EI) 445 (M⁺); mp 247-248 °C (dec.).

Example 17

- 15 2-[4-(3-Cyano-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-yl]-acetamide

- [4-(3-Cyano-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-yl]-acetic acid (Example 15) (48 mg, 0.114 mmol) was treated with Boc₂O (37 mg), NH₄HCO₃ (13 mg) and pyridine (6 µL) in DMF (0.6 mL) at 23 °C for 24 h [*cf. Tetrahedron*
20 *Letters* 1995, 36, 7115]. Obtained as a light yellow solid (14 mg).

MS (ISN) 417 [(M-H)⁻]; mp 250 °C (dec.).

Example 18

3-[8-(2-Methoxy-ethoxy)-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

- 25 Prepared from [2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-5-(2-methoxy-ethoxy)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example O7) (135 mg, 0.251 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a light green solid (82 mg).

- 111 -

MS (EI) 435 (M^+); mp 174-176 °C.

Example 19

4-(3-Cyano-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepine-7-carboxylic acid methyl ester

- 5 Prepared from 5-tert.-butoxycarbonylamino-4-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-2-phenylethynyl-benzoic acid methyl ester (Example O9) (511 mg, 0.95 mmol) by treatment with TFA in CH_2Cl_2 according to the general procedure P. Obtained as an off-white solid (321 mg).

MS (EI) 419 (M^+); mp >250 °C.

10 Example 20

3-(8-Methoxy-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

- Prepared from {2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-5-methoxy-4-phenylethynyl-phenyl}-carbamic acid tert.-butyl ester (Example O8) (359 mg, 0.7 mmol)
15 by treatment with TFA in CH_2Cl_2 according to the general procedure P. Obtained as a yellow-brown solid (87 mg).

MS (EI) 391 (M^+); mp >250 °C.

Example 21

- 3-[8-(2-{2-[2-(2-Methoxy-ethoxy)-ethoxy]-ethoxy}-ethoxy)-4-oxo-7-phenylethynyl-4,5-
20 dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

- Prepared from [2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-5-(2-{2-[2-(2-methoxy-ethoxy)-ethoxy]-ethoxy}-ethoxy)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example O11) (300 mg, 0.437 mmol) by treatment with TFA in CH_2Cl_2 according to the general procedure P. Obtained as a light yellow solid (211 mg).
25 MS (EI) 435 (M^+); mp 140-141 °C (dec.).

Example 22

[4-(3-Cyano-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-yloxy]-acetic acid

Prepared from {5-tert.-butoxycarbonylamino-4-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-2-phenylethynyl-phenoxy}-acetic acid tert.-butyl ester (Example O12) (698 mg, 1.14 mmol) by treatment with TFA in CH₂Cl₂ and anisole according to the general procedure P. Obtained as a yellow solid (265 mg).

- 5 MS (ISN) 434 [(M-H)⁺]; mp 257 °C (dec.).

Example 23

3-(8-Cyanomethyl-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

- 10 Prepared from {5-cyanomethyl-2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-4-phenylethynyl-phenyl}-carbamic acid tert.-butyl ester (Example O13) (266 mg, 0.51 mmol) by treatment with TFA in CH₂Cl₂ and anisole according to the general procedure P. Obtained as a yellow solid (145 mg).

MS (EI) 400 (M⁺); mp 262 °C (dec.).

Example 24

- 15 3-[8-(2,3-Dihydroxy-propoxy)-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

- Prepared from [2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-5-(2,2-dimethyl-[1,3]dioxolan-4-ylmethoxy)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example O16) (265 mg, 0.435 mmol) by treatment with TFA in CH₂Cl₂ and anisole
20 according to the general procedure P. Obtained as a yellow solid (62 mg).

MS (ISP) 452 [(M+H)⁺] and 474 [(M+Na)⁺]; mp 230-234 °C (dec.).

Example 25

4-(3-Iodo-phenyl)-7-(2-methoxy-ethoxy)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- 25 Prepared from [2-[3-(3-iodo-phenyl)-3-oxo-propionylamino]-5-(2-methoxy-ethoxy)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example O15) (1.24 g, 1.89 mmol) by treatment with TFA in CH₂Cl₂ and anisole according to the general procedure P. Obtained as a yellow solid (517 mg).

MS (EI) 536 (M^+); mp 192 °C (dec.).

Example 26

2-[4-(3-Cyano-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-yloxy]-acetamide

- 5 EDC (42 mg, 0.22 mmol) was added to [4-(3-cyano-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-yloxy]-acetic acid (Example 22) (50 mg, 0.11 mmol), NH_4Cl (18 mg, 0.33 mmol) and NMM (56 mg, 0.55 mmol) in DMF (1.1 mL) at 0 °C and the reaction mixture was stirred at 23 °C for 2 h. Obtained as a yellow solid (5 mg).

MS (ISN) 417 [$(M-H)^+$]; mp 250 °C (dec.).

10 Example 27

4-(3-Imidazol-1-yl-phenyl)-7-(2-methoxy-ethoxy)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- Prepared from [[[2-[3-(3-Imidazol-1-yl-phenyl)-3-oxo-propionylamino]-5-(2-methoxy-ethoxy)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example O18) (200 mg, 0.336 mmol) by treatment with TFA in CH_2Cl_2 and anisole according to the general procedure P. Obtained as a yellow solid (28 mg).

MS (EI) 476 (M^+); mp 187-189 °C.

Example 28

- [RS]-3-[4-Oxo-8-(2-oxo-[1,3]dioxolan-4-ylmethoxy)-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from [RS]-[2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-5-(2-oxo-[1,3]dioxolan-4-ylmethoxy)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example O19) (400 mg, 0.67 mmol) by treatment with TFA in CH_2Cl_2 according to the general procedure P. Obtained as a yellow solid (287 mg).

- 25 MS (EI) 477 (M^+); mp 222 °C (dec.).

Example 29

3-(8-Ethoxymethyl-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

Prepared from {2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-5-ethoxymethyl-4-phenylethynyl-phenyl}-carbamic acid tert.-butyl ester (Example O20) (140 mg, 0.26 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (93 mg).

5 MS (EI) 419 (M⁺); mp 229 °C.

Example 30

2,2-Dimethyl-propionic acid 4-(3-cyano-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-ylmethyl ester

Prepared from 2,2-dimethyl-propionic acid 5-tert.-butoxycarbonylamino-4-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-2-phenylethynyl-benzyl ester (Example O21) (156 mg, 0.26 mmol) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a light yellow solid (75 mg).

MS (EI) 475 (M⁺); mp 218 °C.

Example 31

15 3-(8-Hydroxymethyl-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

Prepared from 2,2-dimethyl-propionic acid 4-(3-cyano-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-ylmethyl ester (Example 30) (30 mg, 0.063 mmol) and LiOH.H₂O (8 mg, 0.289 mmol) in THF (2 mL), MeOH (0.4 mL) and H₂O (0.4 mL) at 23 °C for 3 days. Obtained as a yellow solid (17 mg).

MS (EI) 391 (M⁺); mp >255 °C.

Example 32

7-Hydroxymethyl-4-(3-imidazol-1-yl-phenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

25 Prepared from (RS)-[2-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-4-phenylethynyl-5-(tetrahydro-pyran-2-yloxymethyl)-phenyl]-carbamic acid tert.-butyl ester (Example O22) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (77 mg).

MS (EI) 432 (M^+); mp 227 °C.

Example 33

4-(3-Imidazol-1-yl-phenyl)-7-(4-methoxy-piperidin-1-yl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- 5 Prepared from [2-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-5-(4-methoxy-piperidin-1-yl)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example O23) by treatment with TFA in CH_2Cl_2 according to the general procedure P. Obtained as a yellow solid (159 mg).

MS (ISP) 516 [($M+H$) $^+$]; mp 222 °C.

10 Example 34

3-[8-(4-Methoxy-piperidin-1-yl)-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

- Prepared from [2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-5-(4-methoxy-piperidin-1-yl)-4-phenylethynyl-phenyl]-carbamic acid tert.-butyl ester (Example O24) by treatment
15 with TFA in CH_2Cl_2 according to the general procedure P. Obtained as a yellow solid (128 mg).

MS (ISP) 475 [($M+H$) $^+$]; mp 250-251 °C.

Example 35

- [4-(3-Imidazol-1-yl-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-yloxy]-acetonitrile
20

Prepared from {5-cyanomethoxy-2-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-4-phenylethynyl-phenyl}-carbamic acid tert.-butyl ester (Example O25) by treatment with TFA in CH_2Cl_2 according to the general procedure P. Obtained as a yellow solid (43 mg).

MS (EI) 457 (M^+); mp 214 °C.

25 Example 36

3-(8-Cyanomethoxy-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

Prepared from {5-cyanomethoxy-2-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-4-phenylethynyl-phenyl}-carbamic acid tert.-butyl ester (Example O26) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (71 mg).

MS (EI) 416 (M⁺); mp 212 °C.

5 Example 37

8-(4-Fluoro-phenylethynyl)-7-hydroxymethyl-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from (RS)-[4-(4-fluoro-phenylethynyl)-2-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-5-(tetrahydro-pyran-2-yloxymethyl)-phenyl]-carbamic acid tert.-butyl ester (Example O27) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (462 mg).

MS (EI) 450 (M⁺); mp 234 °C (dec.).

Example 38

15 8-(4-Fluoro-phenylethynyl)-7-[(2-hydroxy-ethyl)-methyl-amino]-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from (RS)-[4-(4-fluoro-phenylethynyl)-2-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-5-{methyl-[2-(tetrahydro-pyran-2-yloxy)-ethyl]-amino}-phenyl]-carbamic acid tert.-butyl ester (Example O28) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (73 mg).

20 MS (EI) 493 (M⁺); mp 217 °C (dec.).

Example 39

4-(3-Imidazol-1-yl-phenyl)-7-(4-oxo-piperidin-1-yl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

25 Prepared from {5-(4,4-diethoxy-piperidin-1-yl)-2-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-4-phenylethynyl-phenyl}-carbamic acid tert.-butyl ester (Example O29) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (180 mg).

MS (EI) 499 (M⁺); mp 231 °C (dec.).

Example 40

N-tert.-Butyl-2-[4-(3-cyano-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-yloxy]-acetamide

Prepared from [4-(3-cyano-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-yloxy]-acetic acid (Example 22) (87 mg, 0.2 mmol) by treatment with oxalyl chloride (26 μ L, 0.3 mmol) in DMF (0.6 mL) at 0 °C for 1 h, then with tert.-butylamine (106 μ L, 1.0 mmol) at 0 °C for further 30 min. Obtained as a yellow solid (21 mg).

MS (EI) 490 (M^+); mp >250 °C.

10 Example 41

2-[4-(3-Cyano-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-yloxy]-N-methoxy-acetamide

Prepared from [4-(3-cyano-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-yloxy]-acetic acid (Example 22) (44 mg, 0.1 mmol) by treatment with EDC (38 mg, 0.2 mmol), MeONH₂.HCl (9 mg, 0.11 mmol), NMM (0.021 mL, 0.3 mmol) and HOBT (15 mg, 0.11 mmol) in DMF (1 mL) at 0 to 23 °C for 20 h. Obtained as a yellow solid (36 mg).

MS (ISP) 465 [(M+H)⁺]; mp >250 °C.

Example 42

20 7-(2-Hydroxy-ethoxy)-4-(3-imidazol-1-yl-phenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from (RS)-{2-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-4-phenylethynyl-5-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-phenyl}-carbamic acid tert.-butyl ester (Example O29) by treatment with TFA in CH₂Cl₂ according to the general procedure
25 P. Obtained as a light yellow solid (48 mg).

MS (EI) 462 (M^+); mp 224-227 °C.

Example 43

8-(4-Fluoro-phenyl)-7-(4-hydroxy-piperidin-1-yl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from (RS)-(4'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-2-[4-(tetrahydro-pyran-2-yloxy)-piperidin-1-yl]-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example O31) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a light yellow solid (109 mg).

MS (ISP) 496 [(M+H)⁺]; mp 238-240 °C.

Example 44

10 8-(4-Fluoro-phenylethynyl)-7-(2-hydroxy-ethoxy)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from [5-(2-tert.-butoxy-ethoxy)-4-(4-fluoro-phenylethynyl)-2-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-phenyl]-carbamic acid tert.-butyl ester (Example O32) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a light
15 yellow solid (83 mg).

MS (EI) 480 (M⁺); mp 220-222 °C.

Example 45

8-(4-Fluoro-phenyl)-7-[4-(2-hydroxy-ethoxy)-piperidin-1-yl]-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

20 Prepared from (RS)-(4'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-2-[4-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-piperidin-1-yl]-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example O33) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a light yellow solid (97 mg).

MS (ISP) 540 [(M+H)⁺]; mp 225-227 °C.

25 Example 46

8-(4-Fluoro-phenyl)-7-hydroxymethyl-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from (RS)-[4'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-2-(tetrahydro-pyran-2-yloxymethyl)-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example O34) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a light yellow solid (162 mg).

- 5 MS (EI) 426 [(M)⁺]; mp 180-195 °C.

Example 47

[8-(4-Fluoro-phenyl)-4-(3-imidazol-1-yl-phenyl)-2-oxo-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-yloxy]-acetonitrile

- 10 Prepared from {2-cyanomethoxy-4'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O35) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (11 mg).

MS (EI) 451 (M⁺); mp 164 °C.

Example 48

- 15 7-Dimethylaminomethyl-8-(4-fluoro-phenyl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- 20 Prepared from {2-dimethylaminomethyl-4'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O36) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a brown solid (180 mg).

MS (ISP) 454 (M+H)⁺; mp 115-140 °C (dec.).

Example 49

7-(2,2-Dimethyl-tetrahydro-[1,3]dioxolo[4,5-c]pyrrol-5-yl)-8-(4-fluoro-phenyl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- 25 Prepared from {2-(2,2-dimethyl-tetrahydro-[1,3]dioxolo[4,5-c]pyrrol-5-yl)-4'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O37) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (358 mg).

MS (EI) 537 (M^+); mp 240 °C (dec.).

Example 50

7-(cis-3,4-Dihydroxy-pyrrolidin-1-yl)-8-(4-fluoro-phenyl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- 5 Prepared from 7-(2,2-dimethyl-tetrahydro-[1,3]dioxolo[4,5-c]pyrrol-5-yl)-8-(4-fluoro-phenyl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one (Example 49) (304 mg, 0.57 mmol) by treatment 13% HCl (15 mL) in THF (50 mL) at 23 °C for 16 h. Obtained as a yellow solid (209 mg).

MS (ISP) 498 [$(M+H)^+$]; mp 244 °C.

10 Example 51

8-(4-Fluoro-phenyl)-4-(3-imidazol-1-yl-phenyl)-7-methoxy-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- Prepared from {4'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-2-methoxy-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O38) by treatment with
15 TFA in CH_2Cl_2 according to the general procedure P. Obtained as a yellow solid (182 mg).

MS (EI) 426 (M^+); mp 221 °C (dec.).

Example 52

8-(4-Fluoro-phenyl)-4-(3-imidazol-1-yl-phenyl)-7-(4-oxo-piperidin-1-yl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- 20 Prepared from {2-(1,4-dioxo-8-aza-spiro[4.5]dec-8-yl)-4'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O39) by treatment with TFA in CH_2Cl_2 according to the general procedure P. Obtained as an orange-brown solid (150 mg).

MS (ISP) 494 [$(M+H)^+$]; mp 204 °C.

25 Example 53

8-(4-Fluoro-phenyl)-4-(3-imidazol-1-yl-phenyl)-7-methyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {4'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-2-methyl-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O40) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a light yellow solid (216 mg).

- 5 MS (EI) 410 (M⁺); mp 196 °C.

Example 54

[8-(4-Fluoro-phenyl)-4-(3-imidazol-1-yl-phenyl)-2-oxo-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-yloxy]-acetic acid

- 10 Prepared from {4-tert.-butoxycarbonylamino-4'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-2-yloxy}-acetic acid tert.-butyl ester (Example O41) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a beige solid (570 mg).

MS (ISP) 471 [(M+H)⁺]; mp 209 °C (dec.).

Example 55

- 15 2-[8-(4-Fluoro-phenyl)-4-(3-imidazol-1-yl-phenyl)-2-oxo-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-yloxy]-N-hydroxy-acetamide

- Prepared from [8-(4-fluoro-phenyl)-4-(3-imidazol-1-yl-phenyl)-2-oxo-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-yloxy]-acetic acid (Example 54) (94 mg, 0.2 mmol) by reaction with O-tritylhydroxylamine (61 mg, 0.22 mmol), HOBT (30 mg, 0.22 mmol), N-methylmorpholine (66 µL, 0.6 mmol) and EDC (77 mg, 0.4 mmol) in DMF (2 mL) from 0 to 23 °C for 18 h. After extraction and chromatography the resulting orange solid was stirred with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a light yellow solid (71 mg).

MS (ISP) 486 [(M+H)⁺]; mp 147-157 °C (dec.).

- 25 Example 56

7-Chloro-8-(4-fluoro-phenyl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {2-chloro-4'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O42) by

treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a light yellow solid (35 mg).

MS (EI) 430 (M⁺); mp 209-211 °C.

Example 57

- 5 8-(4-Fluoro-phenyl)-4-(3-imidazol-1-yl-phenyl)-7-(2-methoxy-ethoxy)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from [4'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-2-(2-methoxy-ethoxy)-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example O43) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow
10 solid (96 mg).

MS (EI) 470 (M⁺); mp 196-197 °C.

Example 58

8-(4-Fluoro-phenyl)-7-(2-hydroxy-ethoxy)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- 15 Prepared from {2-(2-tert.-butoxy-ethoxy)-4'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O44) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a light green solid (95 mg).

MS (EI) 456 (M⁺); mp 225 °C.

20 Example 59

8-(4-Fluoro-phenyl)-4-(3-imidazol-1-yl-phenyl)-7-(2-oxo-oxazolidin-3-yl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- Prepared from [4'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-2-(2-oxo-oxazolidin-3-yl)-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example O45) by
25 treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (35 mg).

MS (EI) 481 (M⁺); mp 230 °C.

Example 60

8-(4-Fluoro-2-methyl-phenyl)-4-(3-imidazol-1-yl-phenyl)-7-methoxy-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- Prepared from {4'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-2-methoxy-2'-methyl-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O46) by
5 treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a beige solid (101 mg).

MS (EI) 440 (M⁺); mp 225 °C.

Example 61

- 10 8-(4-Fluoro-phenyl)-7-hydroxy-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- Prepared from {2-tert.-butoxy-4'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O47) by
15 treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (109 mg).

MS (EI) 412 (M⁺); mp 250 °C.

Example 62

8-(2-Fluoro-phenyl)-7-hydroxy-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- 20 Prepared from {2-tert.-butoxy-2'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O48) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (132 mg).

MS (EI) 412 (M⁺); mp 220 °C.

- 25 Example 63

8-(4-Fluoro-phenyl)-7-((R)-3-hydroxy-pyrrolidin-1-yl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from (RS)-[4'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-2-[(R)-3-(tetrahydro-pyran-2-yloxy)-pyrrolidin-1-yl]-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example O49) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (74 mg).

5 MS (EI) 481 (M⁺); mp 155-158 °C.

Example 64

8-(2-Fluoro-phenyl)-4-(3-imidazol-1-yl-phenyl)-7-methoxy-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {2'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-2-methoxy-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O50) by treatment with
10 TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (68 mg).

MS (EI) 426 (M⁺); mp 216 °C (dec.).

Example 65

3-[7-(4-Fluoro-phenyl)-8-hydroxy-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-
15 benzonitrile

Prepared from {2-tert.-butoxy-5-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-4'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O51) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (66 mg).

MS (EI) 371 (M⁺); mp >250 °C.

Example 66

3-[7-(2-Fluoro-phenyl)-8-hydroxy-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from {2-tert.-butoxy-5-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-2'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O52) by treatment with TFA in
25 CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (80 mg).

MS (EI) 371 (M⁺); mp >250 °C.

Example 67

3-[7-(2-Fluoro-phenyl)-8-methoxy-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from {5-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-2'-fluoro-2-methoxy-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O53) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a light yellow solid (51 mg).

MS (EI) 385 (M⁺); mp 245-247 °C.

Example 68

8-(2-Fluoro-phenyl)-7-methoxy-4-[3-(2-methyl-imidazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from (2'-fluoro-2-methoxy-5-[3-[3-(2-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino]-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example O54) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (207 mg).

MS (EI) 440 (M⁺); mp 220-222 °C.

Example 69

5-[7-(2-Fluoro-phenyl)-8-methoxy-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-thiophene-2-carbonitrile

Prepared from {5-[3-(5-cyano-thiophen-2-yl)-3-oxo-propionylamino]-2'-fluoro-2-methoxy-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O55) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (103 mg).

MS (EI) 391 (M⁺); mp >250 °C.

Example 70

8-(2-Fluoro-phenyl)-7-methoxy-4-(3-[1,2,4]triazol-4-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {2'-fluoro-2-methoxy-5-[3-oxo-3-(3-[1,2,4]triazol-4-yl-phenyl)-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O56) by

treatment with TFA in CH_2Cl_2 according to the general procedure P. Obtained as a yellow solid (22 mg).

MS (EI) 427 (M^+); mp 188 °C (dec.).

Example 71

- 5 4-[7-(2-Fluoro-phenyl)-8-methoxy-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-pyridine-2-carbonitrile

Prepared from {5-[3-(2-cyano-pyridin-4-yl)-3-oxo-propionylamino]-2'-fluoro-2-methoxy-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O57) by treatment with TFA in CH_2Cl_2 according to the general procedure P. Obtained as a yellow solid (68 mg).

- 10 MS (EI) 386 (M^+); mp 240-242 °C.

Example 72

8-(4-Fluoro-phenyl)-7-hydroxy-4-[3-(2-methyl-imidazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- 15 Prepared from (2-tert.-butoxy-4'-fluoro-5-{3-[3-(2-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino}-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example O58) by treatment with TFA in CH_2Cl_2 according to the general procedure P. Obtained as a yellow solid (49 mg).

MS (ISP) 427 [$(\text{M}+\text{H})^+$]; mp 260 °C.

Example 73

- 20 4-Fluoro-3-[7-(2-fluoro-phenyl)-8-methoxy-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from {5-[3-(5-cyano-2-fluoro-phenyl)-3-oxo-propionylamino]-2'-fluoro-2-methoxy-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O59) by treatment with TFA in CH_2Cl_2 according to the general procedure P. Obtained as a yellow solid (52 mg).

- 25 MS (ISP) 404 [$(\text{M}+\text{H})^+$]; mp >250 °C.

Example 74

4-[7-(4-Fluoro-phenyl)-8-hydroxy-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-pyridine-2-carbonitrile

Prepared from {2-tert.-butoxy-5-[3-(2-cyano-pyridin-4-yl)-3-oxo-propionylamino]-4'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O60) by treatment with
5 TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (24 mg).

MS (EI) 372 (M⁺); mp 164 °C.

Example 75

8-(2-Fluoro-phenyl)-7-hydroxy-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-
10 benzo[b][1,4]diazepin-2-one

Prepared from {2-tert.-butoxy-2'-fluoro-5-[3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O61) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a light yellow solid (61 mg).

15 MS (ISP) 414 [(M+H)⁺]; mp >250 °C.

Example 76

3-[8-(1,4-Dioxo-8-aza-spiro[4.5]dec-8-yl)-7-(2-fluoro-phenyl)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from {5-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-2-(1,4-dioxo-8-aza-spiro[4.5]dec-8-yl)-2'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O62)
20 by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (132 mg).

MS (ISP) 497 [(M+H)⁺]; mp 253 °C.

Example 77

25 7-(1,4-Dioxo-8-aza-spiro[4.5]dec-8-yl)-8-(2-fluoro-phenyl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {2-(1,4-dioxo-8-aza-spiro[4.5]dec-8-yl)-2'-fluoro-5-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example

O63) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (133 mg).

MS (ISP) 538 [(M+H)⁺]; mp 225 °C.

Example 78

- 5 8-(2-Fluoro-phenyl)-4-(3-imidazol-1-yl-phenyl)-7-(4-oxo-piperidin-1-yl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from 7-(1,4-dioxo-8-aza-spiro[4.5]dec-8-yl)-8-(2-fluoro-phenyl)-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one (Example 77) (54 mg, 0.1 mmol) by stirring in 1N HCl (1 mL) and acetone (1 mL) at 23 °C for 44 h. Obtained as a yellow
10 solid (39 mg).

MS (EI) 493 (M⁺); mp 230 °C.

Example 79

8-(2-Fluoro-phenyl)-7-hydroxy-4-[3-(2-methyl-imidazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- 15 Prepared from (2-tert.-butoxy-2'-fluoro-5-{3-[3-(2-methyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino}-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example O64) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (111 mg).

MS (ISN) 425 [(M-H)⁻]; mp >250 °C.

- 20 Example 80

4-[7-(2-Fluoro-phenyl)-8-hydroxy-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-pyridine-2-carbonitrile

- Prepared from {2-tert.-butoxy-5-[3-(2-cyano-pyridin-4-yl)-3-oxo-propionylamino]-2'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O65) by treatment with
25 TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (47 mg).

MS (ISN) 371 [(M-H)⁻]; mp >250 °C.

Example 81

8-(2-Fluoro-phenyl)-7-hydroxy-4-[3-(2-methylsulfanyl-imidazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from (2-tert.-butoxy-2'-fluoro-5-{3-[3-(2-methylsulfanyl-imidazol-1-yl)-phenyl]-3-oxo-propionylamino}-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example O66) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a light yellow solid (148 mg).

MS (ISN) 457 [(M-H)⁺]; mp >250 °C.

Example 82

10 3-[7-(2,5-Difluoro-phenyl)-8-methoxy-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from {5-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-2',5'-difluoro-2-methoxy-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O67) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (49 mg).

15 MS (EI) 403 (M⁺); mp 251 °C.

Example 83

8-(2,5-Difluoro-phenyl)-7-methoxy-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {2',5'-difluoro-2-methoxy-5-[3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O68) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (78 mg).

MS (EI) 445 (M⁺); mp 241 °C.

Example 84

25 2-[7-(2-Fluoro-phenyl)-8-hydroxy-4-oxo-4,5-dihydro-1H-benzo[b][1,4]diazepin-2-yl]-thiophene-3-carbonitrile

Prepared from {2-tert.-butoxy-5-[3-(3-cyano-thiophen-2-yl)-3-oxo-propionylamino]-2'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O69) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as an orange solid (82 mg).

MS (ISN) 376 [(M-H)⁺]; mp 242 °C.

5 Example 85

5-[7-(2-Fluoro-phenyl)-8-hydroxy-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-thiophene-2-carbonitrile

Prepared from {2-tert.-butoxy-5-[3-(5-cyano-thiophen-2-yl)-3-oxo-propionylamino]-2'-fluoro-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O70) by treatment with
10 TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (126 mg).

MS (EI) 377 (M⁺); mp.

Example 86

8-(4-Fluoro-phenyl)-7-hydroxy-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

15 Prepared from {2-tert.-butoxy-4'-fluoro-5-[3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionylamino]-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O71) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (78 mg).

MS (ISP) 414 [(M+H)⁺]; mp >250 °C.

20 Example 87

3-[7-(2-Fluoro-phenyl)-8-(2-methoxy-ethoxy)-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

Prepared from {5-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-2'-fluoro-2-(2-methoxy-ethoxy)-biphenyl-4-yl}-carbamic acid tert.-butyl ester (Example O72) by treatment with
25 TFA in CH₂Cl₂ according to the general procedure P. Obtained as a light yellow solid (141 mg).

MS (EI) 429 (M⁺); mp 211-213 °C.

Example 88

3-[7-(2-Fluoro-phenyl)-8-hydroxymethyl-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

- Prepared from (RS)-[5-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-2'-fluoro-2-(tetrahydro-pyran-2-yloxymethyl)-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example O73) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a light yellow solid (69 mg).

MS (EI) 385 (M⁺); mp 90-91 °C.

Example 89

- 10 8-(2-Fluoro-phenyl)-7-hydroxy-4-[3-(3-methyl-isoxazol-5-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one

- Prepared from (2'-fluoro-2-(4-methoxy-benzyloxy)-5-{3-[3-(3-methyl-isoxazol-5-yl)-phenyl]-3-oxo-propionylamino}-biphenyl-4-yl)-carbamic acid tert.-butyl ester (Example O74) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (278 mg).

MS (ISP) 428 (M+H)⁺; mp 237 °C.

Example 90

3-[7-(2,5-Difluoro-phenyl)-8-hydroxy-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile

- 20 Prepared from [2-tert.-butoxy-5-[3-(3-cyano-phenyl)-3-oxo-propionylamino]-2',5'-difluoro-biphenyl-4-yl]-carbamic acid tert.-butyl ester (Example O75) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (56 mg).

MS (ISP) 390 (M+H)⁺; mp >250 °C.

Example 91

- 25 8-(4-Fluoro-phenylethynyl)-7-hydroxy-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {5-tert.-butoxy-4-(4-fluoro-phenylethynyl)-2-[3-(3-imidazol-1-yl-phenyl)-3-oxo-propionylamino]-phenyl}-carbamic acid tert.-butyl ester (Example O76) by

treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow solid (55 mg).

MS (EI) 436 (M⁺); mp 247 °C.

Example 92

- 5 8-(4-Fluoro-phenylethynyl)-7-hydroxy-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one

Prepared from {5-tert.-butoxy-4-(4-fluoro-phenylethynyl)-2-[3-oxo-3-(3-[1,2,3]triazol-1-yl-phenyl)-propionylamino]-phenyl}-carbamic acid tert.-butyl ester (Example O77) by treatment with TFA in CH₂Cl₂ according to the general procedure P. Obtained as a yellow
10 solid (56 mg).

MS (EI) 437 (M⁺); mp 243 °C.

The following examples exemplify, that the 4-aryl-8-iodo-1,3-dihydro-benzo[b][1,4]diazepin-2-ones could also serve as starting material for the Sonogashira-coupling as illustrated in synthetic scheme G.

15 Example 93

3-(4-Oxo-7-phenylethynyl-8-thiomorpholin-4-yl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile

Prepared from 3-(7-iodo-4-oxo-8-thiomorpholin-4-yl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile (Example 1) (437 mg, 0.895 mmol) and
20 phenylacetylene (0.15 mL, 1.34 mmol) according to the general procedure K. Obtained as a curry solid (334 mg).

MS (EI) 391 (M⁺); mp 234-235 °C (dec.).

Example 94

- 25 (RS)-3-{4-Oxo-8-(1-oxo-thiomorpholin-4-yl)-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl}-benzonitrile

A mixture of 3-(4-oxo-7-phenylethynyl-8-thiomorpholin-4-yl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile (Example 27) (50 mg, 0.180 mmol) and Davis-

reagent (116 mg, 0.432 mmol) in DCM (4.5 mL) was stirred at 23 °C for 1 h. The product was filtered off and washed with DCM. Obtained as a light yellow solid (16 mg).

MS (ISP) 479 [(M+H)⁺] and 501 [(M+Na)⁺]; mp >250 °C (dec.).

Palladium-catalyzed carbonylation of the 4-(3-iodophenyl)-8-phenylethynyl-1,3-dihydro-
5 benzo[b][1,4]diazepin-2-one in the presence of secondary amines leads directly to the
corresponding amides as shown in synthetic scheme I.

Example 95

3-[8-(2-Methoxy-ethoxy)-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-
2-yl]-benzamide

10 Prepared from 4-(3-iodo-phenyl)-7-(2-methoxy-ethoxy)-8-phenylethynyl-1,3-dihydro-
benzo[b][1,4]diazepin-2-one (Example 25) (268 mg, 0.5 mmol) and hexamethyldisilazane
(0.52 mL, 2.5 mmol) according to the general procedure Q. Obtained as a yellow solid (102
mg).

MS (EI) 453 (M⁺); mp 227-230 °C (dec.).

15 Example I

Tablets of the following composition are produced in a conventional manner:

	<u>mg/Tablet</u>
Active ingredient	100
20 Powdered. lactose	95
White corn starch	35
Polyvinylpyrrolidone	8
Na carboxymethylstarch	10
Magnesium stearate	<u>2</u>
25	Tablet weight 250

Example II

Tablets of the following composition are produced in a conventional manner:

- 134 -

	<u>mg/Tablet</u>
Active ingredient	200
Powdered. lactose	100
White corn starch	64
5 Polyvinylpyrrolidone	12
Na carboxymethylstarch	20
Magnesium stearate	<u>4</u>
Tablet weight	400

Example III


10 Capsules of the following composition are produced:

	<u>mg/Capsule</u>
Active ingredient	50
Crystalline. lactose	60
Microcrystalline cellulose	34
15 Talc	5
Magnesium stearate	<u>1</u>
Capsule fill weight	150

The active ingredient having a suitable particle size, the crystalline lactose and the
20 microcrystalline cellulose are homogeneously mixed with one another, sieved and
thereafter talc and magnesium stearate are admixed. The final mixture is filled into hard
gelatine capsules of suitable size.

Claims

1. Compounds of general formula



I

5 wherein

X is a single bond or an ethynediyl group, wherein,

In case X is a single bond,

R¹ is halogen or phenyl which is optionally substituted with halogen, lower alkyl, halo-lower alkyl, lower alkoxy, halo-lower alkoxy, or cyano;

10 In case X is an ethynediyl group,

R¹ is phenyl, optionally substituted with halogen, lower alkyl, halo-lower alkyl, lower cycloalkyl, lower alkoxy or halo-lower alkoxy;

R² is halogen; hydroxy; lower alkyl; lower halo-alkyl; lower alkoxy; hydroxymethyl; hydroxyethoxy; lower alkoxy-(ethoxy)_n (n = 1 to 4); lower alkoxymethyl; cyanomethoxy;

15 morpholine-4-yl; thiomorpholine-4-yl; 1-oxothiomorpholine-4-yl; 1,1-dioxothiomorpholine-4-yl; 4-oxo-piperidine-1-yl; 4-alkoxy-piperidine-1-yl; 4-hydroxy-piperidine-1-yl; 4-hydroxyethoxy-piperidine-1-yl; 4-lower alkyl-piperazine-1-yl; alkoxycarbonyl; 2-dialkylamino-ethylsulfanyl-; N,N-bis lower alkylamino lower alkyl; carbamoylmethyl; alkylsulfonyl; lower alkoxycarbonyl-lower alkyl; alkylcarboxy-lower

20 alkyl; carboxy-lower alkyl; alkoxycarbonylmethylsulfanyl; carboxymethylsulfanyl; 1,4-dioxo-8-aza-spiro[4.5]dec-8-yl; carboxy-lower alkoxy; cyano-lower alkyl; 2,3-dihydroxy-lower alkoxy; carbamoylmethoxy; 2-oxo-[1,3]-dioxolan-4-yl-lower alkoxy; (2-hydroxy-lower alkyl)-lower alkyl amino; hydroxycarbamoyl-lower alkoxy; 2,2-dimethyl-tetrahydro-[1,3]dioxolo[4,5c]-pyrrol-5-yl; lower alkoxy-carbamoyl-lower alkoxy; 3R-hydroxy-

25 pyrrolidin-1-yl; 3,4-dihydroxy-pyrrolidin-1-yl; 2-oxo-oxazolidin-3-yl; lower alkyl-carbamoylmethoxy; or aminocarbamoyl-lower alkoxy;

R³ is a 5 or 6 membered aryl or heteroaryl which are optionally substituted by

halogen; cyano; nitro; azido; hydroxy; carboxy; morpholine-4-carbonyl; carbamoyl;
thiocarbamoyl; N-hydroxycarbamoyl; trimethylsilyl-ethynyl;

or lower alkyl; lower alkoxy; halo-lower alkyl; 4-lower alkyl-piperazine-1-carbonyl; lower

5 alkylcarbamoyl which are optionally substituted by amino, lower alkylamino, acylamino,
oxo, hydroxy, lower alkoxy, lower alkylthio, or carboxy which is optionally esterified or
amidated;

or an optionally substituted five-membered aromatic heterocycle, which may be optionally
substituted by amino, lower alkylamino, acylamino, oxo, hydroxy, lower alkoxy, lower

10 alkylthio, or carboxy which is optionally esterified or amidated, or lower alkyl which is
optionally substituted by halogen, amino, lower alkylamino, acylamino, hydroxy, lower
alkoxy, lower alkylthio, acyloxy, lower alkenoyl, lower alkylsulfinyl, lower alkylsulfonyl,
cycloalkylsulfinyl, cycloalkylsulfonyl, hydroxyimino, alkoxyimino, carboxy which is
optionally esterified or amidated, lower alkenyl, oxo, cyano, carbamoyloxy, sulfamoyl
15 which is optionally substituted by lower alkyl, or amidino which is optionally substituted
by lower alkyl, -C(NRR')=NR" (where R, R' and R" are hydrogen or lower alkyl)

and their pharmaceutically acceptable addition salts.

2. Compounds according to claim 1, wherein R³ is phenyl substituted in meta position by
cyano; halogen; or imidazolyl which is optionally substituted by lower alkyl or
20 methylsulfonyl; 1,2,3-triazolyl; 1,2,4-triazolyl; or isoxazolyl which is optionally
substituted by lower alkyl.

3. Compounds according to claim 2, being

3-(8-Chloro-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-
benzonitrile;

25 3-[8-(4-Methyl-piperazin-1-yl)-4-oxo-7-phenylethynyl-4,5-dihydro-3H-
benzo[b][1,4]diazepin-2-yl]-benzonitrile;

3-(8-Chloro-4-oxo-7-phenyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-
benzonitrile;

30 [4-(3-Cyano-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-
7-ylsulfonyl]-acetic acid methyl ester;

2-[4-(3-Cyano-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-
benzo[b][1,4]diazepin-7-yl]-acetamide;

3-(8-Methoxy-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-
benzonitrile;

- 3-(8-Cyanomethyl-4-oxo-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl)-benzonitrile;
- 4-(3-Iodo-phenyl)-7-(2-methoxy-ethoxy)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
- 5 4-(3-Imidazol-1-yl-phenyl)-7-(2-methoxy-ethoxy)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
- [RS]-3-[4-Oxo-8-(2-oxo-[1,3]dioxolan-4-ylmethoxy)-7-phenylethynyl-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile;
- 7-Hydroxymethyl-4-(3-imidazol-1-yl-phenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
- 10 [4-(3-Imidazol-1-yl-phenyl)-2-oxo-8-phenylethynyl-2,3-dihydro-1H-benzo[b][1,4]diazepin-7-yl]-acetonitrile;
- 8-(4-Fluoro-phenylethynyl)-7-hydroxymethyl-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
- 15 7-(2-Hydroxy-ethoxy)-4-(3-imidazol-1-yl-phenyl)-8-phenylethynyl-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
- 8-(4-Fluoro-phenyl)-7-[4-(2-hydroxy-ethoxy)-piperidin-1-yl]-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
- 8-(4-Fluoro-phenyl)-7-hydroxy-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
- 20 8-(2-Fluoro-phenyl)-7-methoxy-4-[3-(2-methyl-imidazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
- 8-(2-Fluoro-phenyl)-7-hydroxy-4-[3-(1,2,3-triazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
- 25 8-(2-Fluoro-phenyl)-7-hydroxy-4-[3-(2-methyl-imidazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
- 8-(2-Fluoro-phenyl)-7-hydroxy-4-[3-(2-methylsulfanyl-imidazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
- 8-(2,5-Difluoro-phenyl)-7-methoxy-4-[3-(1,2,3-triazol-1-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
- 30 8-(2-Fluoro-phenyl)-7-hydroxy-4-[3-(3-methyl-isoxazol-5-yl)-phenyl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one;
- 3-[7-(2,5-Difluoro-phenyl)-8-hydroxy-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-benzonitrile;
- 35 8-(4-Fluoro-phenylethynyl)-7-hydroxy-4-(3-imidazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one; and


8-(4-Fluoro-phenylethynyl)-7-hydroxy-4-(3-[1,2,3]triazol-1-yl-phenyl)-1,3-dihydro-benzo[b][1,4]diazepin-2-one.

4. Compounds according to claim 1, wherein R³ is thiophenyl, optionally substituted by cyano or halogen; pyridinyl, optionally substituted in 2-position by cyano or halogen,
 5 or thiazolyl, optionally substituted in 2-position with imidazolyl or 4-methylimidazolyl.
5. Compounds according to claim 4, being
 5-[7-(2-Fluoro-phenyl)-8-methoxy-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-thiophene-2-carbonitrile;
 10 2-[7-(2-Fluoro-phenyl)-8-hydroxy-4-oxo-4,5-dihydro-1H-benzo[b][1,4]diazepin-2-yl]-thiophene-3-carbonitrile;
 4-[7-(2-Fluoro-phenyl)-8-methoxy-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-pyridine-2-carbonitrile;
 4-[7-(4-Fluoro-phenyl)-8-hydroxy-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-pyridine-2-carbonitrile;
 15 4-[7-(2-Fluoro-phenyl)-8-hydroxy-4-oxo-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-pyridine-2-carbonitrile; and
 8-(2-Fluoro-phenyl)-4-[2-(4-methyl-imidazol-1-yl)-thiazol-4-yl]-1,3-dihydro-benzo[b][1,4]diazepin-2-one.
- 20 6. A medicament containing one or more compounds of any one of claims 1 to 5 and pharmaceutically acceptable excipients.
7. A medicament according to claim 6 for the treatment or prevention of acute and/or chronic neurological disorders including psychosis, schizophrenia, Alzheimer's disease, cognitive disorders and memory deficits.
- 25 8. A process for preparing compounds of formula I as defined in claim 1, which process comprises reacting a compound of formula II




with a compound of formula IV or IVa

- 139 -



wherein R is ethyl or buthyl, to a compound of formula III



5

which subsequently undergoes deprotection of the amino group and cyclization, to obtain a compound of formula I of claim 1.

9. A compound according to any one of claims 1 to 5, whenever prepared by a process as claimed in claim 9 or by an equivalent method.
10. The use of a compound according to any one of claims 1 to 5 for the treatment of acute and/or chronic neurological disorders including psychosis, schizophrenia, Alzheimer's disease, cognitive disorders and memory deficits.
11. The use of one or more compounds according to claims 1 to 5 and/or one or more of their pharmaceutically acceptable acid addition salts for the manufacture of medicaments for the treatment or prevention of diseases.
12. The use according to claim 13 for the treatment or prevention of acute and/or chronic neurological disorders including psychosis, schizophrenia, Alzheimer's disease, cognitive disorders and memory deficits.
13. The invention as hereinbefore described.

20
